

## WATER-SOLUBLE RHODAMINE DYE CONJUGATES

This application is a division of pending Application No. 09/661,206, filed on Sept. 14, 2000, which is a division of Application No. 09/433,093, filed on November 3, 1999, now US  
5 Patent No. 6,191,278, issued Feb. 20, 2001, all of which are incorporated herein by reference.

### 1. FIELD OF THE INVENTION

The present invention relates generally to fluorescent dye compounds that are useful as molecular probes. In particular, the present invention relates to fluorescent rhodamine dye  
10 compounds that are photostable and highly water-soluble.

### 2. BACKGROUND OF THE INVENTION

The non-radioactive detection of nucleic acids utilizing fluorescent labels is an important technology in modern molecular biology. By eliminating the need for radioactive labels, safety is  
15 enhanced and the environmental impact and costs associated with reagent disposal is greatly reduced. Examples of methods utilizing such non-radioactive fluorescent detection include automated DNA sequencing, oligonucleotide hybridization methods, detection of polymerase-chain-reaction products, immunoassays, and the like.

In many applications it is advantageous to employ multiple spectrally distinguishable  
20 fluorescent labels in order to achieve independent detection of a plurality of spatially overlapping analytes, i.e., multiplex fluorescent detection. Examples of methods utilizing multiplex fluorescent detection include single-tube multiplex DNA probe assays, PCR, single nucleotide polymorphisms and multi-color automated DNA sequencing. The number of reaction vessels may be reduced thereby simplifying experimental protocols and facilitating the production of  
25 application-specific reagent kits. In the case of multi-color automated DNA sequencing, multiplex fluorescent detection allows for the analysis of multiple nucleotide bases in a single electrophoresis lane thereby increasing throughput over single-color methods and reducing uncertainties associated with inter-lane electrophoretic mobility variations.

Assembling a set of multiple spectrally distinguishable fluorescent labels useful for  
30 multiplex fluorescent detection is problematic. Multiplex fluorescent detection imposes at least six severe constraints on the selection of component fluorescent labels, particularly for

applications requiring a single excitation light source, an electrophoretic separation, and/or treatment with enzymes, e.g., automated DNA sequencing. First, it is difficult to find a set of structurally similar dyes whose emission spectra are spectrally resolved, since the typical emission band half-width for organic fluorescent dyes is about 40-80 nanometers (nm). Second, even if dyes with non-overlapping emission spectra are identified, the set may still not be suitable if the respective fluorescent quantum efficiencies are too low. Third, when several fluorescent dyes are used concurrently, simultaneous excitation becomes difficult because the absorption bands of the dyes are usually widely separated. Fourth, the charge, molecular size, and conformation of the dyes must not adversely affect the electrophoretic mobilities of the analyte. Fifth, the fluorescent dyes must be compatible with the chemistry used to create or manipulate the analyte, e.g., DNA synthesis solvents and reagents, buffers, polymerase enzymes, ligase enzymes, and the like. Sixth, the dye must have sufficient photostability to withstand laser excitation.

Currently available multiple dye sets suitable for use in four-color automated DNA sequencing applications require blue or blue-green laser light to adequately excite fluorescence emissions from all of the dyes making up the set, e.g., argon-ion lasers. As lower cost red lasers become available, a need develops for fluorescent dye compounds and their nucleic acid conjugates which satisfy the above constraints and are excitable by laser light having a wavelength above about 500 nm.

### 3. SUMMARY OF THE INVENTION

These and other objects are furnished by the present invention, which in one aspect provides water-soluble, photostable rhodamine dye compounds that can be used as labels in a variety of biological and non-biological assays. Generally, the rhodamine dye compounds of the invention comprise a rhodamine-type parent xanthene ring substituted at the xanthene C-9 carbon with a substituted phenyl ring. The substituted phenyl ring contains three to five substituents including: an *ortho* carboxyl or sulfonate group; one or more aminopyridinium ("Pyr<sup>+</sup>") groups; and one alkylthio, arylthio or heteroarylthio group. The alkylthio, arylthio or heteroarylthio group is believed to be positioned *para* to the carboxyl or sulfonate group, with the remaining positions being substituted with Pyr<sup>+</sup> groups.

The aminopyridinium groups are attached to the phenyl ring at the pyridinium ring nitrogen

and may be substituted or unsubstituted at the pyridinium ring carbons with one or more of a wide variety of the same or different substituents. The substituents may be virtually any group. However, electron-withdrawing groups (*e.g.*, -NO<sub>2</sub>, -F, -Cl, -CN, -CF<sub>3</sub>, etc.) should not be attached directly to the pyridinium ring carbons, as these substituents may adversely affect the synthesis of the rhodamine dyes. Electron-withdrawing groups may be included on a substituent as long as it is spaced away from the pyridinium ring so as to not adversely affect the synthesis of the dyes. Thus, typical pyridinium ring carbon substituents include, but are not limited to -R, -OR, -SR, -NRR, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R, -C(O)R, -C(O)X, -C(S)R, -C(S)X, -C(O)OR, -C(O)O<sup>-</sup>, -C(S)OR, -C(O)SR, -C(S)SR, -C(O)NRR, -C(S)NRR and -C(NR)NRR, where each R is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, or heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl. The R groups may be further substituted with one or more of the same or different substituents, which are typically selected from the group consisting of -X, -R', =O, -OR', -SR', =S, -NR'R', =NR', -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R', -C(O)R', -C(O)X, -C(S)R', -C(S)X, -C(O)OR', -C(O)O<sup>-</sup>, -C(S)OR', -C(O)SR', -C(S)SR', -C(O)NR'R', -C(S)NR'R' and -C(NR)NR'R', where each X is independently a halogen (preferably -F or -Cl) and each R' is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl or heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl. Preferably, the pyridinium ring carbons are unsubstituted. When substituted, the most preferred substituents are the same or different (C<sub>1</sub>-C<sub>6</sub>) alkyls.

The amino group of the aminopyridinium groups is located at the 4-position of the pyridinium ring. The amino group may be a primary, secondary or tertiary amino group, but is typically a tertiary amino. The nitrogen substituents are typically (C<sub>1</sub>-C<sub>6</sub>) alkyl groups or heteroalkyl groups, and may be the same or different. Alternatively, the nitrogen is substituted with an alkylidyl or heteroalkylidyl bridge having from 2 to 5 backbone atoms such that the substituents and the nitrogen atom taken together form a ring structure, which may be saturated or unsaturated, but is preferably saturated. The bridge substituent may be branched or straight-chain, but is preferably straight-chain, *e.g.*, ethano, propano, butano, etc. The ring structure may contain, in addition to the nitrogen atom of the aminopyridinium, one or more heteroatoms, which are typically selected from the group consisting of O, S and N. When the nitrogen atom is not included in a ring structure, the amino group is preferably dimethylamino. When the nitrogen atom is included in a ring structure, the ring is preferably a morpholino or piperazine ring. Particularly preferred Pyr<sup>+</sup> groups are 4-(dimethylamino)pyridinium, 4-(morpholino) pyridinium, and 1-methyl-4-piperazinyropyridinium.

The alkylthio, arylthio or heteroarylthio group is attached to the phenyl ring via the sulfur

atom and may also be substituted with one or more of the same or different substituents. The nature of the substituents will depend upon whether the group is an alkylthio, arylthio or heteroarylthio. The alkyl chain of an alkylthio group may be substituted with virtually any substituent, including, but not limited to, -X, -R, =O, -OR, -SR, =S, -NRR, =NR, -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO,

5 -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R, -C(O)R, -C(O)X, -C(S)R, -C(S)X, -C(O)OR, -C(O)O<sup>-</sup>, -C(S)OR, -C(O)SR, -C(S)SR, -C(O)NRR, -C(S)NRR and -C(NR)NRR, where each X is independently a halogen (preferably -F or -Cl) and each R is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl or heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl. The R groups may be further substituted with one or more of the same or different substituents, which are typically selected from the group consisting of -X, -R',

10 =O, -OR', -SR', =S, -NR'R', =NR', -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R', -C(O)R', -C(O)X, -C(S)R', -C(S)X, -C(O)OR', -C(O)O<sup>-</sup>, -C(S)OR', -C(O)SR', -C(S)SR', -C(O)NR'R', -C(S)NR'R' and -C(NR)NR'R', where each X is independently a halogen (preferably -F or -Cl) and each R' is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl or heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl.

15 Due to synthetic constraints, when the group is an arylthio or heteroarylthio, the aryl or heteroaryl rings should not be directly substituted with halogens, although halogens may be included in the substituent (*e.g.*, a haloalkyl). Thus, when the group is an arylthio or a heteroarylthio, typical substituents include any of the above-listed alkylthio substituents, with the exception of halogen.

The rhodamine dyes of the invention may include a linker L that can be used to conjugate the

20 dyes, preferably by way of covalent attachment, to other compounds or substances, such as peptides, proteins, antibodies, nucleoside/tides, polynucleotides, polymers, particles, etc. The identity of linker L will depend upon the nature of the desired conjugation. For example, the conjugation may be: (i) mediated by ionic interactions, in which case linker L is a charged group; (ii) mediated by hydrophobic interactions, in which case L is a hydrophobic moiety; (iii) mediated by covalent

25 attachment, in which case L is a reactive functional group (R<sub>x</sub>) that is either capable of forming a covalent linkage with another complementary functional group (F<sub>x</sub>) or is capable of being activated so as to form a covalent linkage with complementary functional group F<sub>x</sub>; or (iv) mediated through the use of pairs of specific binding molecules, such as biotin and avidin/streptavidin, in which case linker L is one member of the pair, *e.g.*, biotin.

30 Linker L is attached to the rhodamine dyes of the invention at the rhodamine-type parent xanthene ring and/or it is included as a substituent on the alkylthio, arylthio or heteroarylthio group



substituting the fully substituted phenyl ring. When linker L is attached to the rhodamine-type parent xanthene ring, it is typically attached to a xanthene nitrogen or at the xanthene C4 carbon. The rhodamine dyes may have multiple linking moieties, but preferably have only a single linking moiety.

Depending upon the particular application, linker L may be attached directly to the rhodamine dye, or indirectly through one or more intervening atoms that serve as a spacer. Linker L can be hydrophobic or hydrophilic, long or short, rigid, semirigid or flexible, depending upon the particular application. When L is positioned at the alkylthio, arylthio or heteroarylthio group, it is preferably attached directly to the molecule. In this latter embodiment, L is a bond.

The new, fully substituted phenyl rings described herein can be used to replace the "bottom ring" or "bottom substituent," *i.e.*, the substituent attached to the xanthene C9 carbon, of virtually any rhodamine dye that is known in the art or that will be later developed. Thus, the new, fully substituted phenyl rings described herein can be covalently attached to the C-9 position of virtually any rhodamine-type parent xanthene ring that is now known or that will be later developed to yield a rhodamine dye without longer absorption and emission maxima and with greater water-solubility.

As the new bottom rings do not deleteriously affect the photostability properties that are characteristic of rhodamine dyes, the new dyes are also highly photostable. Exemplary rhodamine-type parent xanthene rings that can comprise the rhodamine dyes of the invention include, by way of example and not limitation, the xanthene rings ("top rings") of the rhodamine dyes described in U.S. Patent No. 5,936,087; U.S. Patent No. 5,750,409; U.S. Patent No. 5,366,860; U.S. Patent No. 5,231,191; U.S. Patent No. 5,840,999; U.S. Patent No. 5,847,162; U.S. application Serial No. 09/038,191, filed March 10, 1998; U.S. application Serial No. 09/277,793, filed March 27, 1999; U.S. application Serial No. 09/325,243, filed June 3, 1999; PCT Publication WO 97/36960; PCT Publication WO 99/27020; Sauer *et al.*, 1995, J. Fluorescence 5(3):247-261; Arden-Jacob, 1993, *Neue Lanwellige Xanthen-Farbstoffe für Fluoreszenzsonden und Farbstoff Laser*, Verlag Shaker, Germany; and Lee *et al.*, 1992, Nucl. Acids Res. 20(10):2471-2483. Preferred rhodamine-type parent xanthene rings are fluorescent.

In another aspect, the invention provides labeled conjugates comprising a rhodamine dye according to the invention and another molecule or substance. The rhodamine dye is conjugated to the other molecule or substance, typically *via* covalent attachment, through linker L, as previously described. Once conjugated, the rhodamine dye provides a convenient fluorescent label for subsequent detection. The rhodamine dyes of the invention can be used to fluorescently label a wide

variety of molecules and substances, including but not limited to, amino acids, peptides, proteins, antibodies, enzymes, receptors, nucleosides/tides, nucleoside/tide analogs, polynucleotides, polynucleotide analogs, nucleic acids, carbohydrates, lipids, steroids, hormones, vitamins, drugs, metabolites, toxins, organic polymers, etc. The dyes can also be used to label particles such as solid  
5 phase synthesis substrates, nanoparticles, microspheres or liposomes. In embodiments involving nanoparticles, microspheres and/or liposomes, the dye need not include a linking moiety. It can be incorporated into the various particles during their formation. The molecule or substance may be labeled with one or more rhodamine dyes of the invention, which may be the same or different.

In one embodiment, a rhodamine dye of the invention is covalently conjugated to another dye  
10 compound to form an energy-transfer dye pair. The energy-transfer dye pair can be further conjugated to other molecules or substances, as described above, to provide an energy-transfer label. The energy-transfer dye pair generally comprises a donor dye (DD), an acceptor dye (AD), and a linkage linking the donor and acceptor dyes. The donor dye is capable of absorbing light at a first wavelength and emitting excitation energy in response. The acceptor dye is capable of absorbing the  
15 excitation energy emitted by the donor dye and fluorescing at a second wavelength in response thereto. While in many instances the emission wavelength of the donor dye and the excitation wavelength of the acceptor dye will overlap, such overlap is not required. The acceptor dye need only fluoresce in response to the donor dye absorbing light, regardless of the mechanism of action. The linkage serves to facilitate efficient energy transfer between the donor and acceptor dyes.

20 According to this aspect of the invention, at least one of the donor and acceptor dyes is a rhodamine dye according to the invention. Preferably, the acceptor dye is a rhodamine dye according to the invention and the donor dye is a xanthene dye, most typically a fluorescein dye. The exact nature or identity of the donor dye will depend upon the excitation and emission properties of the rhodamine acceptor dye, and will be apparent to those having skill in the art. When covalently conjugated to  
25 enzymatically-incorporatable nucleotides, such as dideoxynucleotide 5'-triphosphates, i.e. "terminators", such energy-transfer dyes are ideally suited for use in sequencing nucleic acids.

Since the rhodamine dyes of the invention may comprise virtually any rhodamine-type parent xanthene ring, the dyes cover a broad range of the visible spectrum, ranging from green to red. Thus, both dyes of an energy-transfer dye pair may be rhodamine dyes of the invention. In this  
30 embodiment, one dye of the invention acts as the donor and another as the acceptor, depending upon their spectral properties.

In another embodiment, a rhodamine dye of the invention, or an energy-transfer dye pair including a rhodamine dye of the invention, is covalently conjugated to a nucleoside/tide, nucleoside/tide analog, polynucleotide or polynucleotide analog to form a labeled conjugate therewith. The dye or dye pair is typically conjugated to the nucleobase moiety of the respective  
5 nucleoside/tide, polynucleotide or analog, but may be conjugated to other portions of the molecule, such as the 5'-terminus, 3'-terminus and/or the phosphate ester internucleoside linkage.

In one preferred embodiment, the labeled conjugate is a labeled polynucleotide or polynucleotide analog that can be used as a primer for generating labeled primer extensions products *via* template-directed enzymatic synthesis reactions. In another preferred embodiment, the labeled  
10 conjugate is a labeled terminator. When used in conjunction with enzymatically-extendable nucleotides or nucleotide or analogs, appropriate polymerizing enzymes and a primed template nucleic acid, such labeled terminators can be used to generate a series of labeled primer extension products *via* template-directed enzymatic synthesis for applications such as nucleic acid sequencing.

In a final aspect, the invention provides methods of using the rhodamine dyes or energy-  
15 transfer dye pairs of the invention to sequence a target nucleic acid. The method generally comprises forming a series of differently-sized primer extension products labeled with a rhodamine dye or energy-transfer dye pair of the invention, separating the series of differently-sized labeled extension products, typically based on size, and detecting the separated labeled extension products based on the fluorescence of the label. The sequence of the target nucleic acid is then assembled according to  
20 known techniques.

The series of differently-sized labeled extension products can be conveniently generated by enzymatically extending a primer-target hybrid according to well-known methods. For example, the series of labeled extension products can be obtained using a primer labeled with a rhodamine dye or dye pair of the invention and enzymatically extending the labeled primer-target hybrid in the presence  
25 of a polymerase, a mixture of enzymatically-extendable nucleotides or nucleotide analogs capable of supporting continuous primer extension (*e.g.*, dATP, dGTP, dCTP and dUTP or dTTP) and at least one, typically unlabeled, terminator that terminates primer extension upon incorporation (*e.g.*, a ddNTP). Alternatively, the series of labeled extension products can be obtained using an unlabeled primer and enzymatically extending the unlabeled primer-target hybrid in the presence of a  
30 polymerase, a mixture of enzymatically-extendable nucleotides or nucleotide analogs capable of supporting continuous primer extension and at least one terminator labeled with a rhodamine dye or

energy-transfer dye pair of the invention. In either embodiment, the polymerase serves to extend the primer with enzymatically-extendable nucleotides or nucleotide analogs until a terminator is incorporated, which terminates the extension reaction. Once terminated, the series of labeled extension products are separated, typically based on size, and the separated labeled extension products detected based on the fluorescence of the labels.

In a particularly advantageous embodiment of this method, a mixture of four different terminators are used in a single extension reaction. Each different terminator is capable of terminating primer extension at a different template nucleotide, *e.g.*, a mixture of 7-deaza-ddATP, ddCTP, 7-deaza-ddGTP and ddTTP or ddUTP, and is labeled with a different, spectrally-resolvable fluorophore, at least one of which is a rhodamine dye or energy-transfer dye pair according to the invention. According to this embodiment, an unlabeled primer-target nucleic acid hybrid is enzymatically extended in the presence of a polymerase, a mixture of enzymatically-extendable nucleotides or nucleotide analogs capable of supporting continuous primer extension and a mixture of the four different labeled terminators. Following separation based on size, a series of separated labeled extension products is obtained in which the emission properties (*i.e.*, color) of each separated extension product reveals the identity of its 3'-terminal nucleotide. In a particularly preferred embodiment, all of the labeled terminators are excitable using a single light source.

Alternatively, terminators may be used in the absence of enzymatically-extendable nucleotides. In this instance, the primer is extended by only a single base. Again, the primer may be labeled, or, alternatively, one or more of the terminators may be labeled. Preferably, a mixture of four different labeled terminators is used, as described above. These "mini sequencing" embodiments are particularly useful for identifying polymorphisms in chromosomal DNA or cDNA.

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 provides single color sequencing data obtained with plasmid pGEM, an unlabeled sequencing primer and the labeled terminator 6-FAM-196-7-deaza-ddATP on an ABI PRISM Model 310 sequencer (PE Biosystems, Foster City, CA).

FIG. 2 provides four-color sequencing data obtained with plasmid pGEM1, an unlabeled sequencing primer and a mixture of four, spectrally resolvable, 3'-fluoro, labeled terminators: 6-FAM-196-7-deaza-ddATP; 5-FAM-dR110-7-deaza-ddGTP; 6-FAM-dJON-ddTTP; and 6-FAM-dROX-ddCTP on an ABI PRISM Model 310 sequencer (PE Biosystems, Foster City, CA).

FIG. 3 provides the fluorescence emission spectra (H<sub>2</sub>O) of four dyes with emission maxima from left to right: **236** (E<sub>max</sub> = 545 nm), **190** (E<sub>max</sub> = 650 nm), **196** (E<sub>max</sub> = 661 nm), **232** (E<sub>max</sub> = 665 nm). The ordinate axis is fluorescence units and the abscissa axis is emission wavelength in nm.

FIG. 4 is a plot of a standard curve of the log v. log graph of the average fluorescence intensity in detection at FL1 (650-685 nm) vs. pg/ml of the IL-8 peptide by a fluorescence-linked immunosorbent assay.

## 5. DETAILED DESCRIPTION OF THE INVENTION

### 5.1 Abbreviations

The abbreviations used throughout the specification to refer to certain nucleobases, nucleosides and/or nucleotides are those commonly employed in the art and are as indicated below:

Expression	Abbreviation
adenine	A
7-deazaadenine	7-deaza-A
N <sup>6</sup> -Δ <sup>2</sup> -isopentenyladenine	6iA
N <sup>6</sup> -Δ <sup>2</sup> -isopentenyl-2-methylthioadenine	2ms6iA
cytosine	C
guanine	G
6-thioguanine	6sG
7-deazaguanine	7-deaza-G
N <sup>2</sup> -dimethylguanine	2dmG
7-methylguanine	7mG
thymine	T
4-thiothymine	4sT
uracil	U
dihydrouracil	D
4-thiouracil	4sU
base Y	Y

Expression	Abbreviation
ribonucleoside-5'-triphosphate	NTP
adenosine-5'-triphosphate	ATP
7-deazaadenosine-5'-triphosphate	7-deaza-ATP
cytidine-5'-triphosphate	CTP
5 guanosine-5'-triphosphate	GTP
7-deazaguanosine-5'-triphosphate	7-deaza-GTP
thymidine-5'-triphosphate	TTP
uridine-5'-triphosphate	UTP
2'-deoxyribonucleoside-5'-triphosphate	dNTP
10 2'-deoxyadenosine-5'-triphosphate	dATP
2'-deoxy-7-deazaadenosine-5'-triphosphate	7-deaza-dATP
2'-deoxycytidine-5'-triphosphate	dCTP
2'-deoxyguanosine-5'-triphosphate	dGTP
2'-deoxy-7-deazaguanosine-5'-triphosphate	7-deaza-dGTP
15 2'-deoxythymidine-5'-triphosphate	dTTP
2'-deoxyuridine-5'-triphosphate	dUTP
2',3'-dideoxyribonucleoside-5'-triphosphate	ddNTP
2',3'-dideoxyadenosine-5'-triphosphate	ddATP
2',3'-dideoxy-7-deazaadenosine-5'-triphosphate	7-deaza-ddATP
20 2',3'-dideoxycytidine-5'-triphosphate	ddCTP
2',3'-dideoxyguanosine-5'-triphosphate	ddGTP
2',3'-dideoxy-7-deazaguanosine-5'-triphosphate	7-deaza-ddGTP
2',3'-dideoxythymidine-5'-triphosphate	ddTTP
2',3'-dideoxyuridine-5'-triphosphate	ddUTP

## 5.2 Definitions

In general, the terms used herein to describe the present invention rely on definitions as

understood and used by those skilled in the art. In particular, chemical structures and substructures are described according to IUPAC recommendations ("Nomenclature of Organic Compounds: A Guide to IUPAC Recommendations 1993, R. Panico, W.H. Powell, and Jean-Claude Richer, Eds., Blackwell Science, Ltd., Oxford, U.K.). As used herein, the following terms are intended to have the following meanings:

"Spectrally Resolvable:" means, in reference to a set of fluorescent dyes and/or energy-transfer dye pairs (collectively referred to herein as "dyes" or "labels"), that the fluorescence emission bands of the respective dyes are sufficiently distinct, *i.e.*, sufficiently non-overlapping, that the dyes, either alone or when conjugated to other molecules or substances, are distinguishable from one another on the basis of their fluorescence signals using standard photodetection systems such as photodetectors employing a series of band pass filters and photomultiplier tubes, charged-coupled devices (CCD), spectrographs, etc., as exemplified by the systems described in U.S. Patent Nos. 4,230,558 and 4,811,218 or in Wheelless *et al.*, 1985, *Flow Cytometry: Instrumentation and Data Analysis*, pp. 21-76, Academic Press, New York. Preferably, all of the dyes comprising a spectrally resolvable set of dyes are excitable by a single light source.

"Nucleobase:" refers to a substituted or unsubstituted nitrogen-containing parent heteroaromatic ring of a type that is commonly found in nucleic acids. Typically, but not necessarily, the nucleobase is capable of forming Watson-Crick and/or Hoogsteen hydrogen bonds with an appropriately complementary nucleobase. Exemplary nucleobases include, but are not limited to, purines such as 2-aminopurine, 2,6-diaminopurine, adenine (A), ethenoadenine, N<sup>6</sup>-Δ<sup>2</sup>-isopentenyladenine (6iA), N<sup>6</sup>-Δ<sup>2</sup>-isopentenyl-2-methylthioadenine (2ms6iA), N<sup>6</sup>-methyladenine, guanine (G), isoguanine, N<sup>2</sup>-dimethylguanine (dmG), 7-methylguanine (7mG), 2-thiopyrimidine, 6-thioguanine (6sG), hypoxanthine and O<sup>6</sup>-methylguanine; 7-deaza-purines such as 7-deazaadenine (7-deaza-A) and 7-deazaguanine (7-deaza-G); pyrimidines such as cytosine (C), 5-propynylcytosine, isocytosine, thymine (T), 4-thiothymine (4sT), 5,6-dihydrothymine, O<sup>4</sup>-methylthymine, uracil (U), 4-thiouracil (4sU) and 5,6-dihydrouracil (dihydrouracil; D); indoles such as nitroindole and 4-methylindole; pyrroles such as nitropyrrole; nebularine; base (Y); etc. Additional exemplary nucleobases can be found in Fasman, 1989, *Practical Handbook of Biochemistry and Molecular Biology*, pp. 385-394, CRC Press, Boca Raton, FL, and the references cited therein. Preferred

nucleobases are purines, 7-deazapurines and pyrimidines. Particularly preferred nucleobases are the normal nucleobases, defined *infra*, and their common analogs, *e.g.*, 2ms6iA, 6iA, 7-deaza-A, D, 2dmG, 7-deaza-G, 7mG, hypoxanthine, 4sT, 4sU and Y.

5                    "Normal Nucleobase:" refers to a nucleobase that is naturally-occurring and encoding, *i.e.*, adenine, cytosine, guanine, thymine or uracil.

                  "Nucleoside:" refers to a compound consisting of a nucleobase covalently linked, typically *via* a heteroaromatic ring nitrogen, to the C1' carbon of a pentose sugar. Typical pentose  
10 sugars include, but are not limited to, those pentoses in which one or more of the carbon atoms are each independently substituted with one or more of the same or different -R, -OR, -NRR or halogen groups, where each R is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl or (C<sub>5</sub>-C<sub>14</sub>) aryl. The pentose sugar may be saturated or unsaturated. Exemplary pentose sugars include, but are not limited to, ribose, 2'-deoxyribose, 2'-(C<sub>1</sub>-C<sub>6</sub>)alkylribose, 2'-(C<sub>1</sub>-C<sub>6</sub>)alkoxyribose, 2'-(C<sub>5</sub>-C<sub>14</sub>)aryloxyribose,  
15 2',3'-dideoxyribose, 2',3'-didehydroribose, 2'-deoxy-3'-haloribose, 2'-deoxy-3'-fluororibose, 2'-deoxy-3'-chlororibose, 2'-deoxy-3'-aminoribose, 2'-deoxy-3'-(C<sub>1</sub>-C<sub>6</sub>)alkylribose, 2'-deoxy-3'-(C<sub>1</sub>-C<sub>6</sub>)alkoxyribose and 2'-deoxy-3'-(C<sub>5</sub>-C<sub>14</sub>)aryloxyribose.

                  When the nucleobase is a purine or a 7-deazapurine, the pentose sugar is attached to the N9-position of the nucleobase. When the nucleobase is a pyrimidine, the pentose sugar is attached to the  
20 N1-position of the nucleobase (*see, e.g.*, Kornberg and Baker, 1992, *DNA Replication*, 2<sup>nd</sup> Ed., Freeman, San Francisco), except for pseudouridine, in which the pentose sugar is attached to the C5 position of the uracil nucleobase. Preferred nucleosides are those in which the nucleobase is a purine, a 7-deazapurine, a pyrimidine, a normal nucleobase or a common analog thereof and the pentose sugar is one of the exemplary pentose sugars listed above.

25                    "Normal Nucleoside:" refers to a compound consisting of a normal nucleobase covalently linked *via* the N1 (C, T or U) or N9 (A or G) position of the nucleobase to the C1' carbon of ribose or 2'-deoxyribose.

30                    "Nucleoside Analog:" refers to a nucleoside in which the pentose sugar is replaced with a pentose sugar analog. Exemplary pentose sugar analogs include, but are not limited to,




substituted or unsubstituted furanoses having more or fewer than 5 ring atoms, *e.g.*, erythroses and hexoses, and substituted or unsubstituted 3-6 carbon acyclic sugars. One or more of the carbon atoms may be independently substituted with one or more of the same or different  $-R$ ,  $-OR$ ,  $-NRR$  or halogen groups, where each  $R$  is independently hydrogen,  $(C_1-C_6)$  alkyl or  $(C_5-C_{14})$  aryl.

5

"Nucleotide:" refers to a nucleoside in which one or more, typically one, of the pentose carbons is substituted with a phosphate ester having the formula:

10




where  $a$  is an integer from 0 to 4. Preferably,  $a$  is 2 and the phosphate ester is attached to the 3'- or 5'-carbon of the pentose. Particularly preferred nucleotides are those which are enzymatically-extendable or enzymatically incorporatable (defined *infra*).

15

"Normal Nucleotide:" refers to a normal nucleoside in which the 3'- or 5'-carbon of the ribose or 2'-deoxyribose sugar is substituted with a phosphate ester having the formula:

20



where  $a$  is an integer from 0 to 2. Preferred normal nucleotides are those in which  $a$  is 2 and the phosphate ester is attached to the 5'-carbon of the ribose (an NTP) or 2'-deoxyribose (a dNTP).

"Nucleotide Analog:" refers to a nucleotide in which the pentose sugar and/or one or more of the phosphate esters is replaced with its respective analog. Exemplary pentose sugar analogs are those previously described in conjunction with nucleoside analogs. Exemplary phosphate ester analogs include, but are not limited to, alkylphosphonates, methylphosphonates, phosphoramidates, 5 phosphotriesters, phosphorothioates, phosphorodithioates, phosphoroselenoates, phosphorodiselenoates, phosphoroanilothioates, phosphoroanilidates, phosphoroamidates, boronophosphates, etc., including any associated counterions, if present.

Also included within the definition of "nucleotide analog" are nucleobase-containing molecules which can be polymerized into polynucleotide analogs in which the DNA/RNA phosphate ester and/or sugar phosphate ester backbone is replaced with a different type of linkage. Such polynucleotide analogs are described in more detail, *infra*.

"Enzymatically-Incorporatable Nucleotide or Nucleotide Analog:" refers to a nucleotide or nucleotide analog which is capable of acting as a substrate for a polymerizing enzyme 15 in a template-directed nucleic acid synthesis reaction such that it is incorporated by the enzyme into a nascent polynucleotide or polynucleotide analog chain. Typical enzymatically-incorporatable nucleotides and nucleotide analogs are those in which the sugar is a pentose. Preferred enzymatically-incorporatable nucleotides are those in which the nucleobase is a purine, a 7-deazapurine, a pyrimidine, a normal nucleobase or a common analog thereof and the pentose sugar is 20 a pentose-5'-triphosphate, such as NTPs, dNTPs and ddNTPs.

"Enzymatically-Extendable Nucleotide or Nucleotide Analog:" refers to an enzymatically-incorporatable nucleotide or nucleotide analog that, once incorporated into the nascent polynucleotide or polynucleotide analog chain, supports incorporation of further nucleotides or 25 nucleotide analogs. Thus, enzymatically-extendable nucleotides or nucleotide analogs have a hydroxyl group that is capable of forming a covalent linkage with another, subsequent nucleotide or nucleotide analog. Typical enzymatically-extendable nucleotides and nucleotide analogs are those in which the sugar is a pentose. Preferred enzymatically-extendable nucleotides are those in which the nucleobase is a purine, a 7-deazapurine, a pyrimidine, a normal nucleobase or a common analog 30 thereof and the pentose sugar is a 3'-hydroxypentose-5'-triphosphate, such as NTPs and dNTPs.

A mixture of enzymatically-extendable nucleotides or nucleotide analogs is said to support

continuous primer extension when the mixture contains an enzymatically-extendable nucleotide or nucleotide analog complementary to each base of the template polynucleotide, *e.g.*, a mixture of dATP, dGTP, dCTP and dUTP or dTTP.



5                   "Terminator:" refers to an enzymatically-incorporatable nucleotide or nucleotide analog that, once incorporated into the nascent polynucleotide chain, terminates further chain extension. Typical terminators are those in which the nucleobase is a purine, a 7-deaza-purine, a pyrimidine, a normal nucleobase or a common analog thereof and the sugar moiety is a pentose which includes a 3'-substituent that blocks further synthesis, such as a ddNTP. Substituents that  
10 block further synthesis include, but are not limited to, amino, deoxy, halogen, alkoxy and aryloxy groups. Exemplary terminators include, but are not limited to, those in which the sugar-phosphate ester moiety is 3'-(C<sub>1</sub>-C<sub>6</sub>)alkylribose-5'-triphosphate, 2'-deoxy- 3'-(C<sub>1</sub>-C<sub>6</sub>)alkylribose-5'-triphosphate, 2'-deoxy-3'-(C<sub>1</sub>-C<sub>6</sub>)alkoxyribose-5'-triphosphate, 2'-deoxy-3'-(C<sub>5</sub>-C<sub>14</sub>)aryloxyribose-5'-triphosphate, 2'-deoxy-3'-haloribose-5'-triphosphate, 2'-deoxy-3'-aminoribose-5'-triphosphate, 2',3'-dideoxyribose-  
15 5'-triphosphate or 2',3'-didehydroribose-5'-triphosphate.

"Nucleoside/tide:" refers to a nucleoside and/or a nucleotide and/or a mixture thereof.

"Polynucleotide:" refers to a linear polymeric chain of nucleoside monomer units that  
20 are covalently connected to one another by phosphate ester internucleoside linkages. Unless stated otherwise, "polynucleotide" as used herein includes polymers of any length, including oligonucleotides, polynucleotides and nucleic acids as those terms are commonly used in the art. Where polynucleotides of specific size ranges are intended, the number of monomer units is specifically delineated. Thus, polynucleotides according to the invention can range in size from a few  
25 monomer units (*e.g.*, 4 to 40), to several hundreds of monomer units, to several thousands of monomer units, or even more monomer units. Whenever a polynucleotide is represented by a sequence of letters, *e.g.* "ATGCCTG," it will be understood that the sequence is presented in the 5'→3' direction. 2'-Deoxyribopolynucleotides are preceded with the letter "d," *e.g.* "d(ATGCCTG)."


Polynucleotides may be comprised of a single type of sugar moiety, as in the case of RNA  
30 and DNA, or mixtures of different sugar moieties, as in the case of RNA/DNA chimeras. Preferred polynucleotides are ribopolynucleotides and 2'-deoxyribopolynucleotides according to the structural

formulae below:



wherein:

- 5 each B is independently a nucleobase, preferably a purine, a 7-deazapurine, a pyrimidine, a normal nucleobase or a common analog thereof;
- each m defines the length of the respective polynucleotide and can range from zero to thousands, to tens of thousands, or even more;
- each R is independently selected from the group consisting of hydrogen, halogen,
- 10  $-R''$ ,  $-OR''$ , and  $-NR''R''$ , where each  $R''$  is independently  $(C_1-C_6)$  alkyl or  $(C_5-C_{14})$  aryl, or two adjacent Rs are taken together to form a bond such that the ribose sugar is 2',3'-didehydroribose; and



, where a is zero, one or two.

In the preferred ribopolynucleotides and 2'-deoxyribopolynucleotides illustrated above, the nucleobases B are covalently attached to the C1' carbon of the sugar moiety as previously described.

"Polynucleotide Analog:" refers to a polynucleotide in which at least one nucleoside monomer unit is a nucleoside analog and/or at least one phosphate ester internucleoside linkage is a phosphate ester analog, as previously defined. Also included within the definition of polynucleotide analogs are polymers in which the phosphate ester and/or sugar phosphate ester internucleoside linkages are replaced with other types of linkages, such as N-(2-aminoethyl)-glycine amides and other amides (*see, e.g.*, Nielsen *et al.*, 1991, Science 254:1497-1500; WO 92/20702; U.S. Patent No. 5,719,262; U.S. Patent No. 5,698,685; morpholinos (*see* U.S. Patent No. 5,698,685; U.S. Patent No. 5,378,841; U.S. Patent No. 5,185,144); carbamates (*see* Stirchak & Summerton, 1987, J. Org. Chem. 52:4202); methylene(methylimino) (*see* Vasseur *et al.*, 1992, J. Am. Chem. Soc. 114:4006); 3'-thioformacetals (*see* Jones *et al.*, 1993, J. Org. Chem. 58:2983); sulfamates (*see* U.S. Patent No. 5,470,967); and others (*see, e.g.*, U.S. Patent No. 5,817,781; Frier & Altman, 1997, Nucl. Acids Res. 25:4429 and the references cited therein).

"Alkyl:" refers to a saturated or unsaturated, branched, straight-chain or cyclic monovalent hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane, alkene or alkyne. Typical alkyl groups include, but are not limited to, methyl; ethyls such as ethanyl, ethenyl, ethynyl; propyls such as propan-1-yl, propan-2-yl, cyclopropan-1-yl, prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl, prop-1-yn-1-yl, prop-2-yn-1-yl, etc.; butyls such as butan-1-yl, butan-2-yl, 2-methyl-propan-1-yl, 2-methyl-propan-2-yl, cyclobutan-1-yl, but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, etc.; and the like. Where specific levels of saturation are intended, the nomenclature "alkanyl," "alkenyl" and/or "alkynyl" is used, as defined below. In preferred embodiments, the alkyl groups are (C<sub>1</sub>-C<sub>6</sub>) alkyl.

"Alkanyl:" refers to a saturated branched, straight-chain or cyclic alkyl radical derived by the removal of one hydrogen atom from a single carbon atom of a parent alkane.

Typical alkanyl groups include, but are not limited to, methanyl; ethanyl; propanyls such as propan-1-yl, propan-2-yl (isopropyl), cyclopropan-1-yl, etc.; butan-1-yl, butan-2-yl (*sec*-butyl), 2-methyl-propan-1-yl (isobutyl), 2-methyl-propan-2-yl (*t*-butyl), cyclobutan-1-yl, etc.; and the like. In preferred embodiments, the alkanyl groups are (C<sub>1</sub>-C<sub>6</sub>)

5 alkanyl.

"Alkenyl:" refers to an unsaturated branched, straight-chain or cyclic alkyl radical having at least one carbon-carbon double bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkene. The radical may be in either the *cis* or *trans* conformation about the double bond(s). Typical alkenyl groups include, but are not limited to, ethenyl; propenyls such as prop-1-en-1-yl, prop-1-en-2-yl, prop-2-en-1-yl, prop-2-en-2-yl, cycloprop-1-en-1-yl; cycloprop-2-en-1-yl; butenyls such as but-1-en-1-yl, but-1-en-2-yl, 2-methyl-prop-1-en-1-yl, but-2-en-1-yl, but-2-en-2-yl, buta-1,3-dien-1-yl, buta-1,3-dien-2-yl, cyclobut-1-en-1-yl, cyclobut-1-en-3-yl, cyclobuta-1,3-dien-1-yl, etc.; and the like. In preferred

10  
15  
embodiments, the alkenyl group is (C<sub>2</sub>-C<sub>6</sub>) alkenyl.

"Alkynyl:" refers to an unsaturated branched, straight-chain or cyclic alkyl radical having at least one carbon-carbon triple bond derived by the removal of one hydrogen atom from a single carbon atom of a parent alkyne. Typical alkynyl groups include, but are not limited to,

20 ethynyl; propynyls such as prop-1-yn-1-yl, prop-2-yn-1-yl, etc.; butynyls such as but-1-yn-1-yl, but-1-yn-3-yl, but-3-yn-1-yl, etc.; and the like. In preferred embodiments, the alkynyl group is (C<sub>2</sub>-C<sub>6</sub>) alkynyl.

"Alkyldiyl:" refers to a saturated or unsaturated, branched, straight-chain or cyclic

25 divalent hydrocarbon radical derived by the removal of one hydrogen atom from each of two different carbon atoms of a parent alkane, alkene or alkyne, or by the removal of two hydrogen atoms from a single carbon atom of a parent alkane, alkene or alkyne. The two monovalent radical centers or each valency of the divalent radical center can form bonds with the same or different atoms. Typical alkyldiyls include, but are not limited to methandiyl; ethyldiyls such as

30 ethan-1,1-diyl, ethan-1,2-diyl, ethen-1,1-diyl, ethen-1,2-diyl; propyldiyls such as propan-1,1-diyl, propan-1,2-diyl, propan-2,2-diyl, propan-1,3-diyl, cyclopropan-1,1-diyl, cyclopropan-1,2-diyl,

prop-1-en-1,1-diyl, prop-1-en-1,2-diyl, prop-2-en-1,2-diyl, prop-1-en-1,3-diyl, cycloprop-1-en-1,2-diyl, cycloprop-2-en-1,2-diyl, cycloprop-2-en-1,1-diyl, prop-1-yn-1,3-diyl, etc.; butyldiyls such as, butan-1,1-diyl, butan-1,2-diyl, butan-1,3-diyl, butan-1,4-diyl, butan-2,2-diyl, 2-methyl-propan-1,1-diyl, 2-methyl-propan-1,2-diyl, cyclobutan-1,1-diyl; cyclobutan-1,2-diyl, 5 cyclobutan-1,3-diyl, but-1-en-1,1-diyl, but-1-en-1,2-diyl, but-1-en-1,3-diyl, but-1-en-1,4-diyl, 2-methyl-prop-1-en-1,1-diyl, 2-methanylidene-propan-1,1-diyl, buta-1,3-dien-1,1-diyl, buta-1,3-dien-1,2-diyl, buta-1,3-dien-1,3-diyl, buta-1,3-dien-1,4-diyl, cyclobut-1-en-1,2-diyl, cyclobut-1-en-1,3-diyl, cyclobut-2-en-1,2-diyl, cyclobuta-1,3-dien-1,2-diyl, cyclobuta-1,3-dien-1,3-diyl, but-1-yn-1,3-diyl, but-1-yn-1,4-diyl, buta-1,3-diyn-1,4-diyl, etc.; and

10 the like. Where specific levels of saturation are intended, the nomenclature alkanyldiyl, alkenyldiyl and/or alkynyldiyl is used. In preferred embodiments, the alkyl diyl group is (C<sub>1</sub>-C<sub>6</sub>) alkyl diyl. Also preferred are saturated acyclic alkanyldiyl radicals in which the radical centers are at the terminal carbons, *e.g.*, methandiyl (methano); ethan-1,2-diyl (ethano); propan-1,3-diyl (propano); butan-1,4-diyl (butano); and the like (also referred to as alkyleneos, defined *infra*).

15 "Alkyleno:" refers to a straight-chain alkyl diyl radical having two terminal monovalent radical centers derived by the removal of one hydrogen atom from each of the two terminal carbon atoms of straight-chain parent alkane, alkene or alkyne. Typical alkyleno groups include, but are not limited to, methano; ethyleneos such as ethano, etheno, ethyno; propyleneos such as propano, prop[1]eno, propa[1,2]dieno, prop[1]yno, etc.; butyleneos such as butano, but[1]eno, but[2]eno, buta[1,3]dieno, but[1]yno, but[2]yno, but[1,3]diyno, etc.; and the like. Where specific levels of saturation are intended, the nomenclature alkano, alkeno and/or alkyno is used. In preferred embodiments, the alkyleno group is (C<sub>1</sub>-C<sub>6</sub>) alkyleno.

25 "Heteroalkyl, Heteroalkanyl, Heteroalkenyl, Heteroalkanyl, Heteroalkyldiyl and Heteroalkyleno:" refer to alkyl, alkanyl, alkenyl, alkynyl, alkyl diyl and alkyleno radicals, respectively, in which one or more of the carbon atoms are each independently replaced with the same or different heteroatomic groups. Typical heteroatomic groups which can be included in these radicals include, but are not limited to, -O-, -S-, -O-O-, -S-S-, -O-S-, -NR'-, =N-N=, -N=N-, 30 -N(O)N-, -N=N-NR'-, -PH-, -P(O)<sub>2</sub>-, -O-P(O)<sub>2</sub>-, -SH<sub>2</sub>-, -S(O)<sub>2</sub>-, -SnH<sub>2</sub>- and the like, where each R' is independently hydrogen, alkyl, alkanyl, alkenyl, alkynyl, aryl, arylaryl, arylalkyl, heteroaryl,

heteroarylalkyl or heteroaryl-heteroaryl as defined herein.

"Acyclic Heteroatomic Bridge:" refers to a divalent bridge in which the backbone atoms are exclusively heteroatoms. Typical acyclic heteroatomic bridges include, but are not limited to, any of the various heteroatomic groups listed above, either alone or in combinations.

"Parent Aromatic Ring System:" refers to an unsaturated cyclic or polycyclic ring system having a conjugated  $\pi$  electron system. Specifically included within the definition of "parent aromatic ring system" are fused ring systems in which one or more of the rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example, indane, indene, phenalene, etc. Typical parent aromatic ring systems include, but are not limited to, aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene, and the like.

"Aryl:" refers to a monovalent aromatic hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. Typical aryl groups include, but are not limited to, radicals derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene, penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene, and the like. In preferred embodiments, the aryl group is ( $C_5$ - $C_{14}$ ) aryl, with ( $C_5$ - $C_{10}$ ) being even more preferred. Particularly preferred aryls are phenyl and naphthyl.

"Aryldiyl:" refers to a divalent aromatic hydrocarbon radical derived by the removal of one hydrogen atom from each of two different carbon atoms of a parent aromatic ring system or by the removal of two hydrogen atoms from a single carbon atom of a parent aromatic ring system. The two monovalent radical centers or each valency of the divalent center can form



bonds with the same or different atom(s). Typical aryldiyl groups include, but are not limited to, divalent radicals derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, coronene, fluoranthene, fluorene, hexacene, hexaphene, hexalene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, octacene, octaphene, octalene, ovalene,


- 5 penta-2,4-diene, pentacene, pentalene, pentaphene, perylene, phenalene, phenanthrene, picene, pleiadene, pyrene, pyranthrene, rubicene, triphenylene, trinaphthalene, and the like. In preferred embodiments, the aryldiyl group is (C<sub>5</sub>-C<sub>14</sub>) aryldiyl, with (C<sub>5</sub>-C<sub>10</sub>) being even more preferred. The most preferred aryldiyl groups are divalent radicals derived from benzene and naphthalene, especially phena-1,4-diyl, naphtha-2,6-diyl and naphtha-2,7-diyl.

10

"Aryleno:" refers to a divalent bridge radical having two adjacent monovalent radical centers derived by the removal of one hydrogen atom from each of two adjacent carbon atoms of a parent aromatic ring system. Attaching an aryleno bridge radical, *e.g.* benzeno, to a parent aromatic ring system, *e.g.* benzene, results in a fused aromatic ring system, *e.g.*

- 15 naphthalene. The bridge is assumed to have the maximum number of non-cumulative double bonds consistent with its attachment to the resultant fused ring system. In order to avoid double-counting carbon atoms, when an aryleno substituent is formed by taking together two adjacent substituents on a structure that includes alternative substituents, the carbon atoms of the aryleno bridge replace the bridging carbon atoms of the structure. As an example, consider the following

20 structure:



wherein:

R<sup>1</sup>, when taken alone is hydrogen, or when taken together with R<sup>2</sup> is (C<sub>5</sub>-C<sub>14</sub>) aryleno; and

- 25 R<sup>2</sup>, when taken alone is hydrogen, or when taken together with R<sup>1</sup> is (C<sub>5</sub>-C<sub>14</sub>) aryleno.

When R<sup>1</sup> and R<sup>2</sup> are each hydrogen, the resultant compound is benzene. When R<sup>1</sup> taken

together with  $R^2$  is  $C_6$  arylene (benzeno), the resultant compound is naphthalene. When  $R^1$  taken together with  $R^2$  is  $C_{10}$  arylene (naphthaleno), the resultant compound is anthracene or phenanthrene. Typical arylene groups include, but are not limited to, aceanthryleno, acenaphthylene, acephenanthryleno, anthraceno, azuleno, benzeno (benzo), chryseno, coroneno, 5 fluorantheno, fluoreno, hexaceno, hexapheno, hexaleno, *as*-indaceno, *s*-indaceno, indeno, naphthaleno (naphtho), octaceno, octapheno, octaleno, ovaleno, penta-2,4-dieno, pentaceno, pentaleno, pentapheno, peryleno, phenaleno, phenanthreno, piceno, pleiadenno, pyreno, pyranthreno, rubiceno, triphenylene, trinaphthaleno, and the like. Where a specific connectivity is intended, the involved bridging carbon atoms (of the arylene bridge) are denoted in brackets, *e.g.*, [1,2]benzeno 10 ([1,2]benzo), [1,2]naphthaleno, [2,3]naphthaleno, etc. Thus, in the above example, when  $R^1$  taken together with  $R^2$  is [2,3]naphthaleno, the resultant compound is anthracene. When  $R^1$  taken together with  $R^2$  is [1,2]naphthaleno, the resultant compound is phenanthrene. In a preferred embodiment, the arylene group is ( $C_5$ - $C_{14}$ ), with ( $C_5$ - $C_{10}$ ) being even more preferred.

15 "Arylaryl:" refers to a monovalent hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a ring system in which two or more identical or non-identical parent aromatic ring systems are joined directly together by a single bond, where the number of such direct ring junctions is one less than the number of parent aromatic ring systems involved. Typical arylaryl groups include, but are not limited to, biphenyl, triphenyl, 20 phenyl-naphthyl, binaphthyl, biphenyl-naphthyl, and the like. When the number of carbon atoms comprising an arylaryl group is specified, the numbers refer to the carbon atoms comprising each parent aromatic ring. For example, ( $C_5$ - $C_{14}$ ) arylaryl is an arylaryl group in which each aromatic ring comprises from 5 to 14 carbons, *e.g.*, biphenyl, triphenyl, binaphthyl, phenyl-naphthyl, etc. Preferably, each parent aromatic ring system of an arylaryl group is independently a ( $C_5$ - $C_{14}$ ) 25 aromatic, more preferably a ( $C_5$ - $C_{10}$ ) aromatic. Also preferred are arylaryl groups in which all of the parent aromatic ring systems are identical, *e.g.*, biphenyl, triphenyl, binaphthyl, trinaphthyl, etc.

"Biaryl:" refers to an arylaryl radical having two identical parent aromatic systems joined directly together by a single bond. Typical biaryl groups include, but are not limited to, 30 biphenyl, binaphthyl, bianthracyl, and the like. Preferably, the aromatic ring systems are ( $C_5$ - $C_{14}$ ) aromatic rings, more preferably ( $C_5$ - $C_{10}$ ) aromatic rings. A particularly preferred biaryl group is

biphenyl.

"Arylalkyl:" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or  $sp^3$  carbon atom, is replaced with an aryl radical.

- 5 Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, 2-phenylethen-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, 2-naphthylethen-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. Where specific alkyl moieties are intended, the nomenclature arylalkanyl, arylakenyl and/or arylalkynyl is used. In preferred embodiments, the arylalkyl group is ( $C_6$ - $C_{20}$ ) arylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl
- 10 group is ( $C_1$ - $C_6$ ) and the aryl moiety is ( $C_5$ - $C_{14}$ ). In particularly preferred embodiments the arylalkyl group is ( $C_6$ - $C_{13}$ ), *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the arylalkyl group is ( $C_1$ - $C_3$ ) and the aryl moiety is ( $C_5$ - $C_{10}$ ).

- "Parent Heteroaromatic Ring System:" refers to a parent aromatic ring system in
- 15 which one or more carbon atoms (and any necessary associated hydrogen atoms) are each independently replaced with the same or different heteroatom. Typical heteratoms to replace the carbon atoms include, but are not limited to, N, P, O, S, Si, etc. Specifically included within the definition of "parent heteroaromatic ring systems" are fused ring systems in which one or more rings are aromatic and one or more of the rings are saturated or unsaturated, such as, for example,
- 20 arsindeole, chromane, chromene, indole, indoline, xanthene, etc. Typical parent heteroaromatic ring systems include, but are not limited to, arsindeole, carbazole,  $\beta$ -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran,
- 25 pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like.

- "Heteroaryl:" refers to a monovalent heteroaromatic radical derived by the removal of one hydrogen atom from a single atom of a parent heteroaromatic ring system. Typical
- 30 heteroaryl groups include, but are not limited to, radicals derived from acridine, arsindeole, carbazole,  $\beta$ -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole,

indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole,


- 5 thiadiazole, thiazole, thiophene, triazole, xanthene, and the like. In preferred embodiments, the heteroaryl group is a 5-14 membered heteroaryl, with 5-10 membered heteroaryl being particularly preferred. The most preferred heteroaryl radicals are those derived from parent heteroaromatic ring systems in which any ring heteroatoms are nitrogens, such as imidazole, indole, indazole, isoindole, naphthyridine, pteridine, isoquinoline, phthalazine, purine, pyrazole, pyrazine, pyridazine, pyridine, pyrrole, quinazoline, quinoline, etc.

"Heteroaryldiyl:" refers to a divalent heteroaromatic radical derived by the removal of one hydrogen atom from each of two different atoms of a parent heteroaromatic ring system or by the removal of two hydrogen atoms from a single atom of a parent heteroaromatic ring system.

- 15 The two monovalent radical centers or each valency of the single divalent center can form bonds with the same or different atom(s). Typical heteroaryldiyl groups include, but are not limited to, divalent radicals derived from acridine, arindole, carbazole,  $\beta$ -carboline, chromane, chromene, cinnoline, furan, imidazole, indazole, indole, indoline, indolizine, isobenzofuran, isochromene, isoindole, isoindoline, isoquinoline, isothiazole, isoxazole, naphthyridine, oxadiazole, oxazole, perimidine, phenanthridine, phenanthroline, phenazine, phthalazine, pteridine, purine, pyran, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolizine, quinazoline, quinoline, quinolizine, quinoxaline, tetrazole, thiadiazole, thiazole, thiophene, triazole, xanthene, and the like. In preferred embodiments, the heteroaryldiyl group is 5-14 membered heteroaryldiyl, with 5-10 membered being particularly preferred. The most preferred heteroaryldiyl groups are divalent radicals derived from parent heteroaromatic ring systems in which any ring heteroatoms are nitrogens, such as imidazole, indole, indazole, isoindole, naphthyridine, pteridine, isoquinoline, phthalazine, purine, pyrazole, pyrazine, pyridazine, pyridine, pyrrole, quinazoline, quinoline, etc.

- "Heteroaryleno:" refers to a divalent bridge radical having two adjacent monovalent radical centers derived by the removal of one hydrogen atom from each of two adjacent atoms of a parent heteroaromatic ring system. Attaching a heteroaryleno bridge radical,

*e.g.* pyridino, to a parent aromatic ring system, *e.g.* benzene, results in a fused heteroaromatic ring system, *e.g.*, quinoline. The bridge is assumed to have the maximum number of non-cumulative double bonds consistent with its attachment to the resultant fused ring system. In order to avoid double-counting ring atoms, when a heteroaryleno substituent is formed by taking together two adjacent substituents on a structure that includes alternative substituents, the ring atoms of the heteroaryleno bridge replace the bridging ring atoms of the structure. As an example, consider the following structure:



wherein:

R<sup>1</sup>, when taken alone is hydrogen, or when taken together with R<sup>2</sup> is 5-14 membered heteroaryleno; and

R<sup>2</sup>, when taken alone is hydrogen, or when taken together with R<sup>1</sup> is 5-14 membered heteroaryleno;

When R<sup>1</sup> and R<sup>2</sup> are each hydrogen, the resultant compound is benzene. When R<sup>1</sup> taken together with R<sup>2</sup> is a 6-membered heteroaryleno (*e.g.*, pyridino), the resultant compound is isoquinoline, quinoline or quinolizine. When R<sup>1</sup> taken together with R<sup>2</sup> is a 10-membered heteroaryleno (*e.g.*, isoquinoline), the resultant compound is, *e.g.*, acridine or phenanthridine. Typical heteroaryleno groups include, but are not limited to, acridino, carbazolo,  $\beta$ -carbolino, chromeno, cinnolino, furano, imidazolo, indazoleno, indoleno, indolizino, isobenzofurano, isochromeno, isoindoleno, isoquinolino, isothiazoleno, isoxazoleno, naphthyridino, oxadiazoleno, oxazoleno, perimidino, phenanthridino, phenanthrolino, phenazino, phthalazino, pteridino, purino, pyrano, pyrazino, pyrazoleno, pyridazino, pyridino, pyrimidino, pyrroleno, pyrrolizino, quinazolino, quinolino, quinolizino, quinoxalino, tetrazoleno, thiadiazoleno, thiazoleno, thiopheno, triazoleno, xantheno, and the like. Where a specific connectivity is intended, the involved bridging atoms (of the heteroaryleno bridge) are denoted in brackets, *e.g.*, [1,2]pyridino, [2,3]pyridino, [3,4]pyridino, etc. Thus, in the above example, when R<sup>1</sup> taken together with R<sup>2</sup> is [1,2]pyridino, the resultant compound is quinolizine. When R<sup>1</sup> taken together with R<sup>2</sup> is [2,3]pyridino, the resultant

compound is quinoline. When R<sup>1</sup> taken together with R<sup>2</sup> is [3,4]pyridino, the resultant compound is isoquinoline. In preferred embodiments, the heteroaryleno group is 5-14 membered heteroaryleno, with 5-10 membered being even more preferred. The most preferred heteroaryleno radicals are those derived from parent heteroaromatic ring systems in which any ring heteroatoms are nitrogens, such as imidazolo, indolo, indazolo, isoindolo, naphthyridino, pteridino, isoquinolino, phthalazino, purino, pyrazolo, pyrazino, pyridazino, pyridino, pyrrolo, quinazolino, quinolino, etc.

"Heteroaryl-Heteroaryl:" refers to a monovalent heteroaromatic radical derived by the removal of one hydrogen atom from a single atom of a ring system in which two or more identical or non-identical parent heteroaromatic ring systems are joined directly together by a single bond, where the number of such direct ring junctions is one less than the number of parent heteroaromatic ring systems involved. Typical heteroaryl-heteroaryl groups include, but are not limited to, bipyridyl, tripyridyl, pyridylpurinyl, bipurinyl, etc. When the number of ring atoms are specified, the numbers refer to the number of atoms comprising each parent heteroaromatic ring systems. For example, 5-14 membered heteroaryl-heteroaryl is a heteroaryl-heteroaryl group in which each parent heteroaromatic ring system comprises from 5 to 14 atoms, *e.g.*, bipyridyl, tripyridyl, etc. Preferably, each parent heteroaromatic ring system is independently a 5-14 membered heteroaromatic, more preferably a 5-10 membered heteroaromatic. Also preferred are heteroaryl-heteroaryl groups in which all of the parent heteroaromatic ring systems are identical. The most preferred heteroaryl-heteroaryl radicals are those in which each heteroaryl group is derived from parent heteroaromatic ring systems in which any ring heteroatoms are nitrogens, such as imidazole, indole, indazole, isoindole, naphthyridine, pteridine, isoquinoline, phthalazine, purine, pyrazole, pyrazine, pyridazine, pyridine, pyrrole, quinazoline, quinoline, etc.

25

"Biheteroaryl:" refers to a heteroaryl-heteroaryl radical having two identical parent heteroaromatic ring systems joined directly together by a single bond. Typical biheteroaryl groups include, but are not limited to, bipyridyl, bipurinyl, biquinolyl, and the like. Preferably, the heteroaromatic ring systems are 5-14 membered heteroaromatic rings, more preferably 5-10 membered heteroaromatic rings. The most preferred biheteroaryl radicals are those in which the heteroaryl groups are derived from a parent heteroaromatic ring system in which any ring


heteroatoms are nitrogens, such as biimidazolyl, biindolyl, biindazolyl, biisoindolyl, binaphthyridinyl, bipteridinyl, biisoquinolinyl, biphtalazinyl, bipurinyl, bipyrazolyl, bipyrazinyl, bipyridazinyl, bipyridinyl, bipyrryl, biquinazolinyl, biquinolyl, etc.

5                    "Heteroarylalkyl:" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or  $sp^3$  carbon atom, is replaced with a heteroaryl radical. Where specific alkyl moieties are intended, the nomenclature heteroarylalkanyl, heteroarylakenyl and/or heterorylalkynyl is used. In preferred embodiments, the heteroarylalkyl group is a 6-20 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety of the heteroarylalkyl is 1-6 membered and the heteroaryl moiety is a 5-14-membered heteroaryl. In particularly preferred embodiments, the heteroarylalkyl is a 6-13 membered heteroarylalkyl, *e.g.*, the alkanyl, alkenyl or alkynyl moiety is 1-3 membered and the heteroaryl moiety is a 5-10 membered heteroaryl.

15                    "Substituted:" refers to a radical in which one or more hydrogen atoms are each independently replaced with the same or different substituent(s). Typical substituents include, but are not limited to,  $-X$ ,  $-R$ ,  $-O^{\cdot}$ ,  $=O$ ,  $-OR$ ,  $-O-OR$ ,  $-SR$ ,  $-S^{\cdot}$ ,  $=S$ ,  $-NRR$ ,  $=NR$ , perhalo ( $C_1-C_6$ ) alkyl,  $-CX_3$ ,  $-CF_3$ ,  $-CN$ ,  $-OCN$ ,  $-SCN$ ,  $-NCO$ ,  $-NCS$ ,  $-NO$ ,  $-NO_2$ ,  $=N_2$ ,  $-N_3$ ,  $-S(O)_2O^{\cdot}$ ,  $-S(O)_2OH$ ,  $-S(O)_2R$ ,  $-C(O)R$ ,  $-C(O)X$ ,  $-C(S)R$ ,  $-C(S)X$ ,  $-C(O)OR$ ,  $-C(O)O^{\cdot}$ ,  $-C(S)OR$ ,  $-C(O)SR$ ,  $-C(S)SR$ ,  $-C(O)NRR$ ,  $-C(S)NRR$  and  $-C(NR)NRR$ , where each  $X$  is independently a halogen (preferably  $-F$  or  $-Cl$ ) and each  $R$  is independently hydrogen, alkyl, alkanyl, alkenyl, alkynyl, aryl, arylalkyl, arylaryl, heteroaryl, heteroarylalkyl or heteroaryl-heteroaryl, as defined herein. The actual substituent substituting any particular group will depend upon the identity of the group being substituted.

25

"Parent Xanthene Ring:" refers to a heteroaromatic ring system of a type typically found in the xanthene class of fluorescent dyes (which includes rhodamine and fluorescein dyes, defined *infra*), *i.e.*, a heteroaromatic ring system having the general structure:



5

In the parent xanthene ring depicted above,  $A^1$  is  $-OH$  or  $-NH_2$  and  $A^2$  is  $=O$  or  $=NH_2^+$ . When  $A^1$  is  $-OH$  and  $A^2$  is  $=O$ , the parent xanthene ring is a fluorescein-type parent xanthene ring, which is defined in more detail, *infra*. When  $A^1$  is  $-NH_2$  and  $A^2$  is  $=NH_2^+$ , the parent xanthene ring is a rhodamine-type parent xanthene ring, which is defined in more detail, *infra*. When  $A^1$  is  $-NH_2$  and

- 10  $A^2$  is  $=O$ , the parent xanthene ring is a rhodol-type parent xanthene ring. In the parent xanthene ring depicted above, one or both nitrogens of  $A^1$  and  $A^2$  (when present) and/or one or more of the carbon atoms at positions C1, C2, C4, C5, C7 and C8, can be independently substituted with a wide variety of the same or different substituents, as is well known in the art. Typical substituents include, but are not limited to,  $-X$ ,  $-R$ ,  $-OR$ ,  $-SR$ ,  $-NRR$ , perhalo ( $C_1-C_6$ ) alkyl,  $-CX_3$ ,  $-CF_3$ ,  $-CN$ ,  
 15  $-OCN$ ,  $-SCN$ ,  $-NCO$ ,  $-NCS$ ,  $-NO$ ,  $-NO_2$ ,  $-N_3$ ,  $-S(O)_2O^-$ ,  $-S(O)_2OH$ ,  $-S(O)_2R$ ,  $-C(O)R$ ,  $-C(O)X$ ,  $-C(S)R$ ,  $-C(S)X$ ,  $-C(O)OR$ ,  $-C(O)O^-$ ,  $-C(S)OR$ ,  $-C(O)SR$ ,  $-C(S)SR$ ,  $-C(O)NRR$ ,  $-C(S)NRR$  and  $-C(NR)NRR$ , where each  $X$  is independently a halogen (preferably  $-F$  or  $-Cl$ ) and each  $R$  is independently hydrogen, ( $C_1-C_6$ ) alkyl, ( $C_1-C_6$ ) alkanyl, ( $C_1-C_6$ ) alkenyl, ( $C_1-C_6$ ) alkynyl, ( $C_5-C_{20}$ ) aryl, ( $C_6-C_{26}$ ) arylalkyl, ( $C_5-C_{20}$ ) arylaryl, heteroaryl, 6-26 membered heteroarylalkyl or 5-20  
 20 membered heteroaryl-heteroaryl. Moreover, the C1 and C2 substituents and/or the C7 and C8 substituents can be taken together to form substituted or unsubstituted ( $C_5-C_{20}$ ) aryleno bridges. Generally, substituents groups which do not tend to quench the fluorescence of the parent xanthene ring are preferred, but in some embodiments quenching substituents may be desirable. Substituents that tend to quench fluorescence of parent xanthene rings are electron-withdrawing groups, such as  
 25  $-NO_2$ ,  $-F$ ,  $-Br$ ,  $-CN$  and  $-CF_3$ .

When  $A^1$  is  $-NH_2$  and/or  $A^2$  is  $=NH_2^+$ , the xanthene nitrogens can be included in bridges involving the same nitrogen atom or adjacent carbon atoms, *e.g.*, ( $C_1-C_{12}$ ) alkylidiyl, ( $C_1-C_{12}$ ) alkyleno, 2-12 membered heteroalkylidiyl and/or 2-12 membered heteroalkyleno bridges.

Any of the substituents substituting carbons C1, C2, C4, C5, C7 or C8 and/or the xanthene




nitrogen atoms (when present) can be further substituted with one or more of the same or different substituents, which are typically selected from the group consisting of -X, -R', =O, -OR', -SR', =S, -NR'R', =NR', -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -NHOH, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R', -P(O)(O<sup>-</sup>)<sub>2</sub>, -P(O)(OH)<sub>2</sub>, -C(O)R', -C(O)X, -C(S)R', -C(S)X, -C(O)OR',  
 5 -C(O)O<sup>-</sup>, -C(S)OR', -C(O)SR', -C(S)SR', -C(O)NR'R', -C(S)NR'R' and -C(NR)NR'R', where each X is independently a halogen (preferably -F or -Cl) and each R' is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, 2-6 membered heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl.

Exemplary parent xanthene rings include, but are not limited to, rhodamine-type parent xanthene rings and fluorescein-type parent xanthene rings, each of which is defined in more detail,

10 *infra*.

"Rhodamine-Type Parent Xanthene Ring:" refers to a parent xanthene ring in which A<sup>1</sup> is -NH<sub>2</sub> and A<sup>2</sup> is =NH<sub>2</sub><sup>+</sup>, *i.e.*, a parent xanthene ring having the general structure:


15



In the rhodamine-type parent xanthene ring depicted above, one or both nitrogens and/or one or  
 20 more of the carbons at positions C1, C2, C4, C5, C7 or C8 can be independently substituted with a wide variety of the same or different substituents, as previously described for the parent xanthene rings. Exemplary rhodamine-type parent xanthene rings include, but are not limited to, the xanthene rings of the rhodamine dyes described in U.S. Patent No. 5,936,087; U.S. Patent No. 5,750,409; U.S. Patent No. 5,366,860; U.S. Patent No. 5,231,191; U.S. Patent No. 5,840,999; U.S.  
 25 Patent No. 5,847,162; U.S. application Serial No. 09/277,793, filed March 27, 1999; PCT Publication WO 97/36960; PCT Publication WO 99/27020; Sauer *et al.*, 1995, J. Fluorescence 5(3):247-261; Arden-Jacob, 1993, *Neue Lanwellige Xanthen-Farbstoffe für Fluoreszenzsonden und Farbstoff Laser*, Verlag Shaker, Germany; and Lee *et al.*, 1992, Nucl. Acids Res. 20(10):2471-

2483. Also included within the definition of "rhodamine-type parent xanthene ring" are the extended-conjugation xanthene rings of the extended rhodamine dyes described in U.S. application Serial No. 09/325,243, filed June 3, 1999.

- 5                    "Fluorescein-Type Parent Xanthene Ring:" refers to a parent xanthene ring in which A<sup>1</sup> is -OH and A<sup>2</sup> is =O, *i.e.*, a parent xanthene ring having the structure:



10

In the fluorescein-type parent xanthene ring depicted above, one or more of the carbons at positions C1, C2, C4, C5, C7 or C8 can be independently substituted with a wide variety of the same or different substituents, as previously described for the parent xanthene rings. Exemplary fluorescein-type parent xanthene rings include, but are not limited to, the xanthene rings of the fluorescein dyes described in U.S. Patent No. 4,439,356; U.S. Patent No. 4,481,136; U.S. Patent No. 5,188,934; U.S. Patent No. 5,654,442; U.S. Patent No. 5,840,999; WO 99/16832; and EP 0 050 684. Also included within the definition of "fluorescein-type parent xanthene ring" are the extended xanthene rings of the fluorescein dyes described in U.S. Patent No. 5,750,409 and U.S. Patent No. 5,066,580.

20

"Xanthene Dye:" refers to a class of fluorescent dyes which consist of a parent xanthene ring substituted at the xanthene C-9 carbon with a substituted phenyl ring or other, typically acyclic, substituent. Common substituted phenyl rings found in xanthene dyes include, *e.g.*, 2-carboxyphenyl, dihalo-2-carboxyphenyl, tetrahalo-2-carboxyphenyl, 2-ethoxycarbonylphenyl, dihalo-2-ethoxycarbonylphenyl and tetrahalo-2-ethoxycarbonylphenyl. Common acyclic substituents found in xanthene dyes include, *e.g.*, carboxyethyl and perfluoroalkyl (*e.g.*, trifluoromethyl, pentafluoroethyl and heptafluoropropyl). Typical xanthene dyes include the fluorescein dyes and the rhodamine dyes, which are described in more detail, *infra*.

"Rhodamine Dye:" refers to the subclass of xanthene dyes in which the xanthene ring is a rhodamine-type parent xanthene ring. Typical rhodamine dyes include, but are not limited to, rhodamine B, 5-carboxyrhodamine, rhodamine X (ROX), 4,7-dichlororhodamine X (dROX), rhodamine 6G (R6G), rhodamine 110 (R110), 4,7-dichlororhodamine 110 (dR110), tetramethyl rhodamine (TAMRA) and 4,7-dichloro-tetramethylrhodamine (dTAMRA). Additional typical rhodamine dyes can be found, for example, in U.S. Patent No. 5,936,087; U.S. Patent No. 5,750,409; U.S. Patent No. 5,366,860; U.S. Patent No. 5,231,191; U.S. Patent No. 5,840,999; U.S. Patent No. 5,847,162; U.S. application Serial No. 09/038,191, filed March 10, 1998; U.S. application Serial No. 09/277,793, filed March 27, 1999; U.S. application Serial No. 09/325,243, filed June 3, 1999; PCT Publication WO 97/36960; PCT Publication WO 99/27020; Sauer *et al.*, 1995, J. Fluorescence 5(3):247-261; Arden-Jacob, 1993, *Neue Lanwellige Xanthen-Farbstoffe für Fluoreszenzsonden und Farbstoff Laser*, Verlag Shaker, Germany; and Lee *et al.*, 1992, Nucl. Acids Res. 20(10):2471-2483.

"Fluorescein Dye:" refers to the subclass of xanthene dyes in which the parent xanthene ring is a fluorescein-type parent xanthene ring. Typical fluorescein dyes include, but are not limited to, 5-carboxyfluorescein (5-FAM), 6-carboxyfluorescein (6-FAM). Additional typical fluorescein dyes can be found, for example, in U.S. Patent No. 5,750,409; U.S. Patent No. 5,066,580; U.S. Patent No. 4,439,356; U.S. Patent No. 4,481,136; U.S. Patent No. 5,188,934; U.S. Patent No. 5,654,442; U.S. Patent No. 5,840,999; PCT publication WO 99/16832; EP 0 050 684; and U.S. application Ser. No. 08/942,067, filed Oct. 1, 1997..

### 5.3 The Rhodamine Dye Compounds

#### 5.3.1 The Compounds Per Se


The rhodamine dyes of the invention are generally rhodamine-type parent xanthene rings substituted at the xanthene C9 position with a new bottom ring. The new bottom ring is a phenyl group which bears five substituents: an *ortho* carboxyl or sulfonate group (or salts thereof); one to three aminopyridinium (Pyr<sup>+</sup>) groups; and one alkylthio, arylthio or heteroarylthio group. The Pyr<sup>+</sup> groups, which may be the same or different, are typically the same and are attached to the new bottom ring *via* the pyridinium ring nitrogen. The alkylthio, arylthio or heteroarylthio group is attached to the new bottom ring *via* the sulfur atom. As previously described in the Summary

section, the aminopyridinium and/or alkylthio, arylthio or heteroarylthio groups may be substituted or unsubstituted. The rhodamine dyes may also include an optional linking moiety, described in more detail, *infra*.

Currently available red-emitting fluorescent dyes, such as rhodamines and cyanines, suffer from undesirable water-solubility and/or photostability characteristics. For example, due to their hydrophobic nature, most commercially-available rhodamine dyes are somewhat insoluble in water. Red-emitting cyanine dyes such as Cy5, although water-soluble, are photo unstable. Thus, available red-emitting rhodamine and cyanine dyes are not well-suited for many aqueous-based biological applications, such as cell staining or nucleic acid sequencing.

By virtue of their new bottom rings, the rhodamine dyes of the invention overcome these limitations. While not intending to be bound by any particular theory, it is believed that the Pyr<sup>+</sup> groups substituting the new bottom ring render the dyes highly water-soluble. Quite importantly, rhodamine dyes substituted with these new bottom rings retain their characteristic photostability. Moreover, the new bottom ring tends to shift the emissions spectral properties of the dyes to the red by about 5-30 nm, as compared with corresponding rhodamines dyes comprising a conventional bottom ring. Thus, the rhodamine dyes of the invention are ideally suited for use as water-soluble laser dyes and in aqueous-based biological applications such as cell staining and nucleic acid sequencing. Quite significantly, when used to label terminators in nucleic acid sequencing applications, the new rhodamine dyes of the invention do not effect obscuring impurities which electrophoretically migrate in the range of DNA sequencing fragments. Obscuring impurities are commonly observed with conventionally-labeled terminators and are generally thought to be caused by unincorporated labelled-terminators. As a consequence, sequencing data obtained with terminators labeled with the new rhodamine dyes of the invention is typically much higher in quality than that obtained with conventional terminators. Sequencing reactions obtained with terminators labeled with the new rhodamine dyes of the present invention may require less sample purification, or "clean up" than those employing conventional terminators.

The rhodamine dyes of the invention are generally compounds according to structural formula (I):



5 including any associated counterions, wherein:

E is carboxylic acid, sulfonic acid, or a salt thereof;

Y is a rhodamine-type parent xanthene ring connected to the illustrated fully substituted phenyl ring at the C-9 carbon;

each  $\text{Pyr}^+$  is independently a substituted or unsubstituted aminopyridinium group  
 10 connected to the illustrated phenyl ring *via* the pyridinium ring nitrogen;

S is sulfur;

n is one, two, or three; and

Z a substituted or unsubstituted ( $\text{C}_1\text{-C}_{12}$ ) alkyl, ( $\text{C}_5\text{-C}_{14}$ ) aryl or heteroaryl.

The invention is based, in part, on the discovery that replacing the ring or substituent  
 15 attached to the xanthene C-9 carbon of conventional rhodamine dyes with the new bottom rings described herein yields rhodamine dyes having superior water-solubility, photo stability and/or excitation and emission spectral properties. As a consequence, those of skill in the art will recognized that in the rhodamine dyes of structural formula (I), rhodamine-type parent xanthene ring Y can be derived from virtually any fluorescent rhodamine dye that is now known or that will  
 20 be later developed. Exemplary rhodamine-type parent xanthene rings that can comprise Y include, but are not limited to, any of the substituted or unsubstituted xanthene rings of the rhodamine dyes described in U.S. Patent No. 5,936,087; U.S. Patent No. 5,750,409; U.S. Patent No. 5,366,860; U.S. Patent No. 5,231,191; U.S. Patent No. 5,840,999; U.S. Patent No. 5,847,162; U.S. application Serial No. 09/038,191, filed March 10, 1998; U.S. application Serial No. 09/277,793,  
 25 filed March 27, 1999; U.S. application Serial No. 09/325,243, filed June 3, 1999; PCT Publication WO 97/36960; PCT Publication WO 99/27020; Sauer *et al.*, 1995, J. Fluorescence 5(3):247-261; Arden-Jacob, 1993, *Neue Lanwellige Wanthen-Farbstoffe für Fluoreszenzsonden und Farbstoff Laser*, Verlag Shaker, Germany; and Lee *et al.*, 1992, Nucl. Acids Res. 20(10):2471-2483, the

disclosures of which are incorporated herein by reference.

While the dyes of structural formula (I) are useful in virtually any aqueous-based applications employing red-emitting fluorescent dyes, such as, for example, as water-soluble laser dyes, rhodamines according to structural formula (I) which incorporate one or more optional

5 linking moieties are particularly useful, as they can be specifically and/or permanently conjugated to other compounds and/or substances so as to label the compounds or substances for subsequent detection. The linking moieties are attached to the rhodamine-type parent xanthene ring, either at a xanthene nitrogen, the xanthene C4 carbon, and/or the Z substituent. Thus, preferred rhodamine dyes of the invention are compounds according to structural formula (I) in which the C4 carbon

10 atom of Y, one or both nitrogen atoms of Y and/or substituent Z are substituted with a linking moiety of the formula -L-, wherein L is a bond or linker. When the dye includes multiple linkers L, each may be the same or different.

The nature of linker L will depend upon the particular application, point of attachment and type of conjugation desired. Linker L may be attached directly to the dye, or it may be spaced away

15 from the dye through one or more intervening atoms that serve as a linker. In the former embodiment, L represents a bond. In the latter embodiment, L represents a linker of more than one atom. The linker can be hydrophilic or hydrophobic, long or short, rigid, semirigid or flexible, depending upon the particular application. The linker can be optionally substituted with one or more substituents or one or more additional linking groups, which may be the same or different,

20 thereby providing a "polyvalent" linking moiety capable of conjugating with multiple molecules or substances. Preferably, however, linker L does not include such additional substituents or linking groups.

A wide variety of linkers L comprised of stable bonds are known in the art, and include by way of example and not limitation, alkylidiyls, substituted alkylidiyls, alkylenos, substituted

25 alkylenos, heteroalkylidiyls, substituted heteroalkylidiyls, heteroalkylenos, substituted heteroalkylenos, acyclic heteroatomic bridges, aryldiyls, substituted aryldiyls, arylaryldiyls, substituted arylaryldiyls, arylalkylidiyls, substituted arylalkylidiyls, heteroaryldiyls, substituted heteroaryldiyls, heteroaryl-heteroaryldiyls, substituted heteroaryl-heteroaryldiyls, heteroarylalkylidiyls, substituted heteroarylalkylidiyls, heteroaryl-heteroalkylidiyls, substituted




30 heteroaryl-heteroalkylidiyls, and the like. Thus, linker L may include single, double, triple or aromatic carbon-carbon bonds, nitrogen-nitrogen bonds, carbon-nitrogen, carbon-oxygen bonds


and/or carbon-sulfur bonds, and may therefore include functionalities such as carbonyls, ethers, thioethers, carboxamides, sulfonamides, ureas, urethanes, hydrazines, etc. In one embodiment, linker L has from 1-20 non-hydrogen atoms selected from the group consisting of C, N, O, and S and is composed of any combination of ether, thioether, amine, ester, carboxamide, sulfonamides, 5 hydrazide, aromatic and heteroaromatic bonds.

Choosing a linker having properties suitable for a particular application is within the capabilities of those having skill in the art. For example, where a rigid linker is desired, L may be a rigid polypeptide such as polyproline, a rigid polyunsaturated alkylidiyl or an arylidiyl, biaryldiyl, arylaryldiyl, arylalkylidiyl, heteroaryldiyl, biheteroaryldiyl, heteroarylalkylidiyl,

10 heteroaryl-heteroaryldiyl, etc. Where a flexible linker is desired, L may be a flexible polypeptide such as polyglycine or a flexible saturated alkanyldiyl or heteroalkanyldiyl. Hydrophilic linkers may be, for example, polyalcohols or polyethers such as polyalkyleneglycols. Hydrophobic linkers may be, for example, alkylidiyls or arylidiyls.

Linkers suitable for use in most biological applications include (C<sub>1</sub>-C<sub>12</sub>) alkylidiyls,

15 particularly alkanylenes such as methano (-CH<sub>2</sub>-), ethano (-CH<sub>2</sub>-CH<sub>2</sub>-), propano (-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-), butano (-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-), pentano (-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-) and hexano (-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-); (C<sub>5</sub>-C<sub>20</sub>) arylidiyls, particularly phenyl-1,3-diyl () , phenyl-1,4-diyl () , naphthalen-2,6-diyl () and naphthalen-2,7-diyl

() ); and (C<sub>6</sub>-C<sub>26</sub>) arylalkylidiyls, particularly those having the structural formula

20  $-(CH_2)_i-\phi$  or  $-(CH_2)_i-\psi$ , where each  $i$  is independently an integer from 1 to 6,  $\phi$  is phenyldiyl (especially phenyl-1,3-diyl or phenyl-1,4-diyl) and  $\psi$  is naphthyldiyl (especially naphthalen-2,6-diyl or naphthalen-2,7-diyl). Rigid linkers that are suitable for attaching the dyes of the invention to one another or to other dyes to create energy-transfer dye pairs (described in more detail, *infra*) include  $-(CH_2)_i-NR''-C(O)-\phi$  and  $-(CH_2)_i-NR''-C(O)-\psi$ , where  $i$ ,  $R''$ ,  $\phi$  and  $\psi$  are as previously defined.

25 Particularly preferred are those linkers in which  $i$  is 1. Analogs of all of these linkers L containing one or more heteroatomic groups, particularly those selected from the group consisting of O, S, N and NR'', where R'' is hydrogen or (C<sub>1</sub>-C<sub>6</sub>) alkyl, can also be conveniently used to space linkers from the rhodamine dyes of the invention. Linkers L tailored to specific applications are discussed in more detail, *infra*.

Rhodamine dyes including a linking moiety can be conjugated to a variety of different molecules and substances using a plethora of different conjugation means. For example, the conjugation can be mediated *via* hydrophobic interactions, ionic attraction, through the use of pairs of specific binding molecules such as biotin and avidin/streptavidin or through covalent

5 attachment. When conjugation *via* hydrophobic interactions is desired, linker L is a hydrophobic moiety that is capable of forming hydrophobic interactions with a hydrophobic moiety on the molecule or substance to be conjugated. Typical hydrophobic moieties include, but are not limited to, unsubstituted and substituted aryl, arylalkyl, arylaryl, heteroaryl, heteroarylalkyl and heteroaryl-heteroaryl groups. When the hydrophobic moiety is substituted, the substituents are  
10 preferably nonpolar, more preferably hydrophobic. Suitable hydrophobic moieties for forming non-covalent conjugates will be apparent to those of skill in the art.

When conjugation *via* ionic attraction is desired, L is a charged moiety having a net charge of a polarity opposite to a net charge on the molecule or substance to be conjugated. Typical charged moieties include, by way of example and not limitation, quaternary ammoniums,  
15 carboxylates and sulfonates, including salts thereof. A variety of cyclic quaternary ammoniums that are suitable for use in linkers are described in U.S. Patent No. 5,863,753 (*see, e.g.*, Cols. 8-9), the disclosure of which is incorporated herein by reference.

When conjugation *via* pairs of specific binding molecules such as biotin and avidin/streptavidin is desired, L will constitute one member of the binding pair. The molecule or  
20 substance to be conjugated will bear the other member of the binding pair. Where one of the members of the specific binding pair is a small molecule, such as biotin or a hormone, that member preferably comprises L. A variety of biotins capable of being covalently linked to reactive functional groups such as amines are commercially available (*e.g.*, Molecular Probes, Eugene, OR). These biotins can be incorporated into the dyes of the invention to yield biotin-labeled dyes  
25 suitable for non-covalent conjugation to a variety of avidin/streptavidin-labeled molecules or substances.

Preferably, L is capable of mediating conjugation *via* covalent attachment. In this preferred embodiment, L bears a reactive functional group ( $R_x$ ). Covalent conjugates are obtained by reacting a rhodamine dye of the invention including a reactive group  $R_x$  with a molecule or  
30 substance that contains, or is modified to contain, one or more functional groups  $F_x$  that are complementary to reactive group  $R_x$ .



The exact identities of  $R_x$  and  $F_x$  will depend upon the nature of the desired covalent linkage and the chemistry used to form the covalent linkage. Generally, reactive group  $R_x$  is a functional group that is capable of reacting with a complementary functional group  $F_x$  under specified reaction conditions to form a covalent linkage. However, those of skill in the art will recognize that a variety of functional groups that are typically unreactive under certain reaction conditions can be activated to become reactive. Groups that can be activated to become reactive include, *e.g.*, carboxylic acids and esters, including salts thereof. Such groups are referred to herein as "activatable precursors" and are specifically intended to be included within the expression "reactive group."

Pairs of reactive groups  $R_x$  and complementary groups  $F_x$  suitable for forming covalent linkages with one another under a variety of different reaction conditions are well-known. Any of these complementary pairs of groups can be used to covalently conjugate the rhodamine dyes of the invention to other compounds or substances. In one convenient embodiment, reactive group  $R_x$  and complementary functional group  $F_x$  comprise complementary electrophiles and nucleophiles (or their respective activatable precursors). In another convenient embodiment, reactive group  $R_x$  is a photoactivatable group that becomes chemically reactive only after illumination with light of an appropriate wavelength and complementary functional group  $F_x$  is a group capable forming a covalent linkage with the chemically reactive species. Such photoactivatable groups can be conveniently used to photo cross-link the rhodamine dyes of the invention to other molecules and/or substances.

As understood in the art, "activated esters" generally have the formula  $-C(O)\Omega$ , where  $\Omega$  is a good leaving group. Exemplary good leaving groups include, by way of example and not limitation: oxysuccinimidyl; N-succinimidyl; oxysulfosuccinimidyl; 1-oxybenzotriazolyl; and  $-OR^a$ , where  $R^a$  is selected from the group consisting of ( $C_4$ - $C_{20}$ ) cycloalkyl (*e.g.*, cyclohexyl), heterocycloalkyl, ( $C_5$ - $C_{20}$ ) aryl, ( $C_5$ - $C_{20}$ ) aryl substituted with one or more of the same or different electron-withdrawing groups (*e.g.*,  $-\text{NO}_2$ ,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{CN}$ ,  $-\text{CF}_3$ , etc.), heteroaryl, and heteroaryl substituted with one or more of the same or different electron-withdrawing groups, *n*-dialkylaminoalkyls (*e.g.*, 3-dimethylaminopropyl) and N-morpholinomethyl, or  $R^a$  is used to form an anhydride of the formula  $-\text{OCOR}^b$  or  $-\text{OCNR}^b\text{NHR}^c$ , where  $R^b$  and  $R^c$  are each independently selected from the group consisting of ( $C_1$ - $C_6$ ) alkyl, ( $C_1$ - $C_6$ ) perhaloalkyl, ( $C_1$ - $C_6$ ) perfluoroalkyl and ( $C_1$ - $C_6$ ) alkoxy. A preferred activated ester is NHS ester.

Exemplary photoactivatable groups suitable for conjugation *via* light-activated cross-linking include, but are not limited to, azido ( $-N_3$ ), 4-azido-phenyl and 2-nitro-4-azido-phenyl. Conjugation using photoactivatable groups typically involves illuminating a mixture comprising the photoactivatable dyes and the molecule or substance to be conjugated, followed by separation of unreacted dyes and byproducts.

As will be recognized by those of skill in the art, reactive group  $R_x$  can comprise any electrophilic, nucleophilic or photoactivatable groups. The selection of reactive group  $R_x$  used to covalently conjugate the rhodamine dyes of the invention to the other molecule or substance typically depends upon the identity of the complementary functional group  $F_x$  on the molecule or substance to be conjugated. The types of complementary functional groups typically present on molecules or substances to be conjugated include, but are not limited to, amines, thiols, alcohols, phenols, aldehydes, ketones, phosphates, imidazoles, hydrazines, hydroxylamines, mono- and disubstituted amines, halides, epoxides, sulfonate esters, carboxylic acids or carboxylates, or a combination of these groups. A single type of complementary functional group may be available on the molecule or substance (which is typical for polysaccharides), or a variety of different complementary functional groups may be available (*e.g.* amines, thiols, alcohols, phenols), which is typical for proteins. The molecule or substance may be conjugated to more than one rhodamine dye, which may be the same or different. Although some selectivity can be obtained by carefully controlling the reaction conditions, selectivity of conjugation is best obtained by appropriate choice of reactive group  $R_x$  in light of the available complementary functional group(s)  $F_x$ . In instances where the molecule or substance to be conjugated does not contain available complementary functional group(s)  $F_x$ , it can be modified to contain such groups using any of a variety of standard techniques.


In a preferred embodiment, reactive group  $R_x$  is a group that reacts with, or that can be readily activated to react with, an amine, a thiol or an alcohol. In a particularly preferred embodiment, one of reactive group  $R_x$  or complementary functional group  $F_x$  is a carboxylic acid (or a salt thereof) or an activated ester, most preferably a N-hydroxysuccinimidyl (NHS) ester, and the other is an amine, preferably a primary amine. The NHS ester may be conveniently obtained by reacting a rhodamine dye of the invention including a carboxylic acid reactive group  $R_x$  with N-hydroxysuccinimide in the presence of an activating agent (*e.g.*, dicyclohexylcarbodiimide) according to known methods.

For a discussion of the various reactive groups  $R_x$  and respective complementary functional groups  $F_x$  that can be conveniently used to covalently conjugate the rhodamine dyes of the invention to a variety of biological and other molecules or substances, as well as reaction conditions under which the conjugation reactions can be carried out, *see* Haugland, 1996,

5 *Molecular Probes Handbook of Fluorescent Probes and Research Chemicals*, Molecular Probes, Inc.; Brinkley, 1992, *Bioconjugate Chem.* 3:2 and Garman, 1997, *Non-Radioactive Labelling: A Practical Approach*, Academic Press, London, as well as the references cited in all of the above.

In one illustrative embodiment, rhodamine dyes of the invention which are capable of being covalently conjugated to other compounds and/or substances are compounds according to

10 structural formula (Ia):



including any associated counter ions, wherein:

15 E, Y,  $\text{Pyr}^+$  and S are as previously defined for structural formula (I);


$Z^1$  is a substituted or unsubstituted ( $\text{C}_1\text{-C}_{12}$ ) alkylidyl, ( $\text{C}_5\text{-C}_{14}$ ) or heteroarylidy;

L is a bond or a linker as previously described;

$R_x$  is a reactive group as previously described; and

n is one, two, or three.

20 In another illustrative embodiment, rhodamine dyes of the invention which are capable of being covalently conjugated to other compounds and/or substances are compounds according to structural formula (Ib):



5

including any associated counterions, wherein:

E, Y,  $\text{Pyr}^+$ , S and Z are as previously defined for structural formula (I);

L and  $\text{R}_x$  are as previously defined for structural formula (Ia), where L is connected  
10 to the C4 carbon atom or a nitrogen atom of Y; and

n is one, two, or three..

While not intending to be bound by any particular theory, it is believed that the  $\text{Pyr}^+$  groups on the new bottom rings in the compounds of structural formulae (I), (Ia) and (Ib) account for the water-solubility of the rhodamine dyes of the invention.

15 As will be illustrated more thoroughly below, the  $\text{Pyr}^+$  groups are introduced into the rhodamine dyes of the invention by displacing fluorine atoms of the corresponding tetrafluororhodamine precursor with the desired aminopyridine reactant, e.g. Scheme (I), using a reaction similar to that described in Weiss, R. *etal*, 1995, *Angew. Chem. Int. Ed. Engl.* 34:1319-21; and Koch, A. *etal*, 1993, *Jour. Org. Chem.* 58:1409-14. As a consequence, unless mixtures of  
20 different aminopyridine reactants are used in the displacement reaction, all of the  $\text{Pyr}^+$  groups on the new bottom phenyl rings are identical. According to the reaction, the  $\text{Pyr}^+$  groups are attached to the new bottom phenyl ring *via* the pyridinium ring nitrogen.

The pyridinium ring carbons may be independently substituted with a wide variety of the same or different substituents. The desired substituents are introduced as substituents on the  
25 aminopyridine reactants. These carbons may be substituted with virtually any group, with one caveat: in order to avoid deleteriously affecting the displacement reaction, the pyridine ring

carbons should not be directly substituted with electron-withdrawing groups, *e.g.*, -F, -Cl, -CF<sub>3</sub>, -NO<sub>2</sub>, -CN, -N<sub>3</sub>, etc. However, such electron-withdrawing groups can be included on the substituents, as long as it is not attached directly to a pyridine ring carbon. For example, while a pyridine ring carbon should not be directly substituted with -F, -Cl or -CF<sub>3</sub>, it may be substituted

- 5 with other, less electronegative or electron-withdrawing, haloalkyls (*e.g.*, -CH<sub>2</sub>-CH<sub>2</sub>F). Identifying substituents which are suitably non-electron-withdrawing is within the capabilities of those having skill in the art. Typical groups useful for substituting the pyridinium ring carbons include, but are not limited to, -R, -OR, -SR, -NRR, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R, -C(O)R, -C(O)X, -C(S)R, -C(S)X, -C(O)OR, -C(O)O<sup>-</sup>, -C(S)OR, -C(O)SR, -C(S)SR, -C(O)NRR, -C(S)NRR and
- 10 -C(NR)NRR, where each R is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl or heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl. The R groups may be further substituted with one or more of the same or different substituents, which are typically selected from the group consisting of -X, -R', =O, -OR', -SR', =S, -NR'R', =NR', -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R', -C(O)R', -C(O)X, -C(S)R', -C(S)X, -C(O)OR', -C(O)O<sup>-</sup>, -C(S)OR', -C(O)SR', -C(S)SR',
- 15 -C(O)NR'R', -C(S)NR'R' and -C(NR)NR'R', where each X is independently a halogen (preferably -F or -Cl) and each R' is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, 2-6 membered heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl. Preferably, the pyridinium ring carbons are unsubstituted. When substituted, the most preferred substituents are the same or different (C<sub>1</sub>-C<sub>6</sub>) alkyls. Preferably, the pyridinium carbons are unsubstituted. When substituted, the substituents are preferably the same
- 20 or different (C<sub>1</sub>-C<sub>6</sub>) alkyls.

The amino groups of the Pyr<sup>+</sup> substituents may be a primary, secondary or tertiary amino group, but is typically a tertiary amino. The nitrogen substituents, R, are typically the same or different (C<sub>1</sub>-C<sub>6</sub>) alkyl or heteroalkyl. The alkyl or heteroalkyl can be further substituted with one or more of the same or different groups, as previously described for R, above.

- 25 Alternatively, the nitrogen atom may be included in a ring structure having from 2 to 5 ring atoms. The ring may contain, in addition to the amino nitrogen atom, one or more of the same or different heteroatoms, which are typically selected from the group consisting of O, S and N. The ring atoms can be further substituted with any of the previously described substituent groups. Preferably, the amino group is dimethylamino or morpholino.


- 30 In the compounds of structural formulae (I), (Ia) and (Ib), the Z or Z' substituent may also be substituted or unsubstituted, but is preferably unsubstituted. When Z is an alkyl or Z' is an

alkyldiyl, virtually any group can be used to substitute Z or Z<sup>1</sup>. Typical substituents include, but are not limited to, -X, -R, =O, -OR, -SR, =S, -NRR, =NR, -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R, -C(O)R, -C(O)X, -C(S)R, -C(S)X, -C(O)OR, -C(O)O<sup>-</sup>, -C(S)OR, -C(O)SR, -C(S)SR, -C(O)NRR, -C(S)NRR and -C(NR)NRR, where each X is  
 5 independently a halogen (preferably -F or -Cl) and each R is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, 2-6 membered heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl. The R groups may be further substituted with one or more of the same or different substituents, which are typically selected from the group consisting of -X, -R', =O, -OR', -SR', =S, -NR'R', =NR', -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R', -C(O)R', -C(O)X, -C(S)R', -C(S)X, -C(O)OR',  
 10 -C(O)O<sup>-</sup>, -C(S)OR', -C(O)SR', -C(S)SR', -C(O)NRR', -C(S)NRR' and -C(NR)NRR', where each X is independently a halogen (preferably -F or -Cl) and each R' is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, 2-6 membered heteroalkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl or heteroaryl.

However, referring to Scheme (I), it has been observed that the displacement reaction does not proceed efficiently with arylthio and heteroarylthio compounds **108** that are substituted at the  
 15 aromatic ring atoms with halogen groups. Thus, when Z is an aryl or heteroaryl and/or Z<sup>1</sup> is an aryldiyl or heteroaryldiyl, the aromatic ring atoms should not be substituted directly with halogens. However, as described above for the pyridinium ring carbon substituents, the halogens can be included in a substituent (*e.g.*, a haloalkyl), so long as the substituent is not sufficiently  
 electronegative to disrupt the displacement reaction. Thus, with the exception of halogen, groups  
 20 to substitute aryl, heteroaryl, aryldiyl and heteroaryldiyl Z and Z<sup>1</sup> groups are typically any of the groups described above for when Z is an alkyl or Z<sup>1</sup> is an alkyldiyl.

The rhodamine dyes of the invention will now be more fully described by reference to various preferred embodiments. In one preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formulae (I), (Ia) and (Ib) in which each Pyr<sup>+</sup> is  
 25 the same and has the structure:

(P.1)



5

including any associated counter ions, wherein:

$R^{22}$  is selected from the group consisting of hydrogen and  $(C_1-C_6)$  alkyl;

$R^{23}$  is selected from the group consisting of hydrogen and  $(C_1-C_6)$  alkyl;

10  $R^{24}$ , when taken alone, is selected from the group consisting of  $(C_1-C_6)$  alkyl, or when taken together with  $R^{24'}$  is  $(C_4-C_{10})$  alkylidyl,  $(C_4-C_6)$  alkyleno, heteroalkylidyl or heteroalkyleno;

$R^{24'}$ , when taken alone, is selected from the group consisting of  $(C_1-C_6)$  alkyl, or when taken together with  $R^{24}$  is  $(C_4-C_{10})$  alkylidyl,  $(C_4-C_6)$  alkyleno, heteroalkylidyl or heteroalkyleno;

15  $R^{25}$  is selected from the group consisting of hydrogen and  $(C_1-C_6)$  alkyl; and

$R^{26}$  is selected from the group consisting of hydrogen and  $(C_1-C_6)$  alkyl.

In the  $\text{Pyr}^+$  substituents of structural formula (P.1), the dashed line at the pyridinium ring nitrogen indicates the point of attachment to the phenyl ring in the compounds of structural

20 formulae (I), (Ia) and/or (Ib). Preferred  $\text{Pyr}^+$  substituents according to structural formula (P.1) are those in which  $R^{22}$ ,  $R^{23}$ ,  $R^{25}$  and  $R^{26}$  are each hydrogen and/or in which  $R^{24}$  and  $R^{24'}$ , when taken alone, are each the same  $(C_1-C_6)$  alkyl or, when taken together are a 4-6 membered heteroalkyleno having a single oxygen heteroatom. Particularly preferred  $\text{Pyr}^+$  substituents according to structural formula (P.1) are 4-(dimethylamino)pyridinium and 4-(morpholino)pyridinium.

25 In another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formulae (I) and (Ib) in which Z is selected from the group consisting of  $(C_1-C_{12})$  alkyl,  $(C_1-C_{12})$  alkyl substituted with one or more of the same or different  $W^1$  groups;  $(C_5-C_{14})$  aryl,  $(C_5-C_{14})$  aryl substituted with one or more of the same or different  $W^2$  groups, 5-14

membered heteroaryl or heteroaryl independently substituted with one or more of the same or different  $W^2$  groups, wherein:

each  $W^1$  is independently selected from the group consisting of -X, -R, =O, -OR, -SR, =S, -NRR, =NR, -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, =N<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>,  
 5 -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R, -C(O)R, -C(O)X, -C(S)R, -C(S)X, -C(O)OR, -C(O)O<sup>-</sup>, -C(S)OR, -C(O)SR, -C(S)SR, -C(O)NRR, -C(S)NRR and -C(NR)NRR, where each X is independently a halogen (preferably -F or -Cl) and each R is independently hydrogen or (C<sub>1</sub>-C<sub>6</sub>) alkyl; and

each  $W^2$  is independently selected from the group consisting of -R, -OR, -SR, -NRR, -CX<sub>3</sub>, -CN, -OCN, -SCN, -NCO, -NCS, -NO, -NO<sub>2</sub>, -N<sub>3</sub>, -S(O)<sub>2</sub>O<sup>-</sup>, -S(O)<sub>2</sub>OH, -S(O)<sub>2</sub>R,  
 10 -C(O)R, -C(O)X, -C(S)R, -C(S)X, -C(O)OR, -C(O)O<sup>-</sup>, -C(S)OR, -C(O)SR, -C(S)SR, -C(O)NRR, -C(S)NRR and -C(NR)NRR, R is as previously defined for  $W^1$ .

Preferably, Z is unsubstituted. However, when Z is a substituted (C<sub>5</sub>-C<sub>14</sub>) aryl or a substituted heteroaryl, the most preferred substituents are those that are not electron-withdrawing. The most preferred heteroaryl groups, whether substituted or unsubstituted, are those in which any  
 15 heteroatoms are nitrogens. Especially preferred amongst these preferred heteroaryls are pyridinyl and purinyl. The most preferred aryl groups, whether substituted or unsubstituted are phenyl and naphthyl.


In another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ia) in which Z<sup>1</sup> is selected from the group consisting of (C<sub>1</sub>-C<sub>12</sub>)  
 20 alkylidiyl, (C<sub>1</sub>-C<sub>12</sub>) alkylidiyl substituted with one or more of the same or different  $W^1$  groups; (C<sub>5</sub>-C<sub>14</sub>) arylidiyl, (C<sub>5</sub>-C<sub>14</sub>) arylidiyl substituted with one or more of the same or different  $W^2$  groups, 5-14 membered heteroaryldiyl or heteroaryldiyl independently substituted with one or more of the same or different  $W^2$  groups, wherein  $W^1$  and  $W^2$  are as defined above.

Preferably, Z<sup>1</sup> is unsubstituted. However, when Z<sup>1</sup> is a substituted (C<sub>5</sub>-C<sub>14</sub>) arylidiyl or a  
 25 substituted heteroaryldiyl, the most preferred substituents are those that are not electron-withdrawing. The most preferred heteroaryldiyl groups, whether substituted or unsubstituted, are those in which any heteroatoms are nitrogens. Especially preferred amongst these preferred heteroaryldiyls are pyridindiyl and purindiyl. The most preferred arylidiyl groups, whether substituted or unsubstituted are phenydiyl and naphthadiyl, especially phena-1,3-diyl, phena-1,4-  
 30 diyl, naphtha-2,6-diyl and naphtha-2,7-diyl.

In yet another preferred embodiment, the rhodamine dyes of the invention are compounds



according to structural formula (Ia) in which rhodamine-type parent xanthene ring is a compound according to structural formula (Y-1):



including any associated counterions, wherein:

R<sup>1</sup> and R<sup>2</sup> are each independently selected from the group consisting of hydrogen and (C<sub>1</sub>-C<sub>6</sub>) alkyl;

10 R<sup>3</sup>, when taken alone, is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl and (C<sub>5</sub>-C<sub>14</sub>) arylaryl, or when taken together with R<sup>3'</sup> is (C<sub>4</sub>-C<sub>6</sub>) alkylidyl or (C<sub>4</sub>-C<sub>6</sub>) alkyleno, or when taken together with R<sup>2</sup> or R<sup>4</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;

R<sup>3'</sup>, when taken alone, is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl and (C<sub>5</sub>-C<sub>14</sub>) arylaryl, or when taken together with R<sup>3</sup> is (C<sub>4</sub>-C<sub>6</sub>) alkylidyl or (C<sub>4</sub>-C<sub>6</sub>)

15 alkyleno, or when taken together with R<sup>2</sup> or R<sup>4</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;

R<sup>4</sup>, when taken alone, is selected from the group consisting of hydrogen and (C<sub>1</sub>-C<sub>6</sub>) alkyl, or when taken together with R<sup>3</sup> or R<sup>3'</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;

R<sup>5</sup>, when taken alone, is selected from the group consisting of hydrogen and (C<sub>1</sub>-C<sub>6</sub>) alkyl, or when taken together with R<sup>6</sup> or R<sup>6'</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;

20 R<sup>6</sup>, when taken alone, is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl and (C<sub>5</sub>-C<sub>14</sub>) arylaryl, or when taken together with R<sup>6'</sup> is (C<sub>4</sub>-C<sub>6</sub>) alkylidyl or (C<sub>4</sub>-C<sub>6</sub>) alkyleno, or when taken together with R<sup>5</sup> or R<sup>7</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;

R<sup>6'</sup>, when taken alone, is selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>5</sub>-C<sub>14</sub>) aryl and (C<sub>5</sub>-C<sub>14</sub>) arylaryl, or when taken together with R<sup>6</sup> is (C<sub>4</sub>-C<sub>6</sub>) alkylidyl or (C<sub>4</sub>-C<sub>6</sub>)

25 alkyleno, or when taken together with R<sup>5</sup> or R<sup>7</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;




R<sup>7</sup>, when taken alone, is selected from the group consisting of hydrogen and (C<sub>1</sub>-C<sub>6</sub>) alkyl, or when taken together with R<sup>6</sup> or R<sup>6'</sup> is (C<sub>2</sub>-C<sub>6</sub>) alkylidyl or (C<sub>2</sub>-C<sub>6</sub>) alkyleno;

R<sup>8</sup>, when taken alone, is selected from the group consisting of hydrogen and (C<sub>1</sub>-C<sub>6</sub>) alkyl;

and

$R^9$  indicates the point of attachment to phenyl bottom ring.

In another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ia) in which rhodamine-type parent xanthene ring is a compound according to structural formula (Y-2), (Y-3) or (Y-4):



including any associated counterions, wherein:

$R^1$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^8$  and  $R^9$  are as previously defined for structural formula (Y-1);

and

$R^{10}$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ ,  $R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  are each independently

selected from the group consisting of hydrogen and (C<sub>1</sub>-C<sub>6</sub>) alkyl, or R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup> and R<sup>13</sup> taken together are (C<sub>5</sub>-C<sub>14</sub>) aryleno or (C<sub>5</sub>-C<sub>14</sub>) aryleno substituted with one or more of the same or different (C<sub>1</sub>-C<sub>6</sub>) alkyl, or R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> taken together are (C<sub>5</sub>-C<sub>14</sub>) aryleno or (C<sub>5</sub>-C<sub>14</sub>) aryleno substituted with one or more of the same or different (C<sub>1</sub>-C<sub>6</sub>) alkyl.

5 The dashed bonds in structural formulae (Y-2) and (Y-4) indicate bonds which can each, independently of one another, be a single or a double bond. When these bonds are double bonds, one of R<sup>10</sup> or R<sup>11</sup> and one of R<sup>12</sup> or R<sup>13</sup> are taken together to form a bond and one of R<sup>18</sup> or R<sup>19</sup> and one of R<sup>20</sup> or R<sup>21</sup> are taken together to form a bond. When these bond are single bonds, the R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> substituents are as defined above.

10 In still another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ib) in which rhodamine-type parent xanthene ring Y is selected from the group consisting of (Y-1), (Y-2), (Y-3) and (Y-4), where R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> are as previously defined, and either R<sup>3'</sup> or R<sup>4</sup> indicates the point of attachment of substituent L. When substituent L is attached to R<sup>3'</sup>, R<sup>4</sup> is as  
15 previously defined. When substituent L is attached to R<sup>4</sup>, R<sup>3'</sup> is as previously defined.

In still another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formulae (Ia) and (Ib) in which Y is selected from the group consisting of (Y-1), (Y-2), (Y-3) and (Y-4), and further in which:

- any alkyl groups are alkanyls selected from the group consisting of methanyl,  
20 ethanyl and propanyl;
- any aryl groups are phenyl or naphthyl;
- any arylaryl groups are biphenyl;
- any alkyldiyl or alkyleno bridges formed by taking R<sup>2</sup> together with R<sup>3</sup> or R<sup>3'</sup> are alkanyldiyls or alkanos, especially those selected from the group consisting of ethano, propano,  
25 1,1-dimethylethano, 1,1-dimethylpropano and 1,1,3-trimethylpropano;
- any alkyldiyl or alkyleno bridges formed by taking R<sup>3</sup> together with R<sup>3'</sup> are alkanyldiyl or alkano, especially butano;
- any alkyldiyl or alkyleno bridges formed by taking R<sup>4</sup> together with and R<sup>3</sup> or R<sup>3'</sup> are alkanyldiyls or alkanos, especially those selected from the group consisting of ethano, propano,  
30 1,1-dimethylethano, 1,1-dimethylpropano and 1,1,3-trimethylpropano;
- any alkyldiyl or alkyleno bridges formed by taking R<sup>5</sup> together with R<sup>6</sup> or R<sup>6'</sup> are

alkanyldiyls or alkanos, especially those selected from the group consisting of ethano, propano, 1,1-dimethylethano, 1,1-dimethylpropano and 1,1,3-trimethylpropano;

any alkylidiyl or alkyleno bridges formed by taking  $R^6$  together with  $R^6$  are alkanyldiyl or alkanos, especially butano;






- 5 any alkylidiyl or alkyleno bridges formed by taking  $R^7$  together with  $R^6$  or  $R^6$  are alkanyldiyls or alkanos, especially those selected from the group consisting of ethano, propano, 1,1-dimethylethano, 1,1-dimethylpropano and 1,1,3-trimethylpropano;






any aryleno bridges formed by taking  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$  and  $R^{13}$  together are benzo; and





any aryleno bridges formed by taking  $R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  together are benzo.

- 10 In yet another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ia) in which rhodamine-type parent xanthene ring Y is selected from the following group of compounds, where  $R^9$  is as previously defined for structural formula (Y-1):

15 (Y-20a)	
(Y-21a)	
20 (Y-22a)	

	(Y-23a)	
5	(Y-24a)	
	(Y-25a)	
10	(Y-31a)	
15	(Y-34a)	

	(Y-35a)	
5	(Y-36a)	
	(Y-39a)	
10	(Y-41a)	
15	(Y-42a)	

(Y-43a)	
5 (Y-44a)	
(Y-45a)	
10 (Y-46a)	

15

In yet another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ia) in which:

Y is selected from the group consisting of (Y-1), (Y-2), (Y-3), (Y-4), (Y-20a), (Y-21a), (Y-22a), (Y-23a), (Y-24a), (Y-25a), (Y-31a), (Y-34a), (Y-35a), (Y-36a), (Y-39a), (Y-41a),

(Y-42a), (Y-43a), (Y-44a), (Y-45a), and (Y-46a);

Z<sup>1</sup> is selected from the group consisting of (C<sub>1</sub>-C<sub>12</sub>) alkylene, (C<sub>1</sub>-C<sub>12</sub>) alkano, (C<sub>5</sub>-C<sub>10</sub>) aryl, phenyl, phenyl, phena-1,4-diyl, naphthadiyl, naphtha-2,6-diyl and heteroaryl, pyridindyl and purindyl.

5 In yet another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ia) in which:

L is a bond; and

Y is selected from the group consisting of (Y-1), (Y-2), (Y-3), (Y-4), (Y-20a), (Y-21a), (Y-22a), (Y-23a), (Y-24a), (Y-25a), (Y-31a), (Y-34a), (Y-35a), (Y-36a), (Y-39a), (Y-41a),  
10 (Y-42a), (Y-43a), (Y-44a), (Y-45a), and (Y-46a);

In still another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ia) in which:

R<sub>x</sub> is any of the electrophilic or nucleophilic groups; and

Y is selected from the group consisting of (Y-1), (Y-2), (Y-3), (Y-4), (Y-20a), (Y-21a), (Y-22a), (Y-23a), (Y-24a), (Y-25a), (Y-31a), (Y-34a), (Y-35a), (Y-36a), (Y-39a), (Y-41a),  
15 (Y-42a), (Y-43a), (Y-44a), (Y-45a), and (Y-46a);

In yet another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ib) in which Y is selected from the group consisting of (Y-1), (Y-2), (Y-3), (Y-4), (Y-20a), (Y-21a), (Y-22a), (Y-23a), (Y-24a), (Y-25a), (Y-31a), (Y-34a), (Y-35a),  
20 (Y-36a), (Y-39a), (Y-41a), (Y-43a) and (Y-46a), where substituent L is attached to the xanthene C4 carbon. In structures (Y-23a), (Y-24a) and (Y-25a), the C4 carbon in the illustrated tautomer is bridged to the xanthene nitrogen. Those of skill in the art will recognize that due to tautomerism, the carbons labeled C4 and C5 (in the definition of rhodamine-type parent xanthene ring) are essentially equivalent. Thus, in the illustrated structures of compounds (Y-23a), (Y-24a) and (Y-25a), substituent L is bonded to the C5 carbon. Particularly preferred compounds according to this  
25 aspect of the invention are those compounds in which:

Z is selected from the group consisting of (C<sub>1</sub>-C<sub>12</sub>) alkyl, (C<sub>1</sub>-C<sub>12</sub>) alkanyl, (C<sub>5</sub>-C<sub>10</sub>) aryl, phenyl, naphthyl, naphth-1-yl, naphth-2-yl, pyridyl and purinyl; and/or

L is selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>1</sub>-C<sub>6</sub>) alkano, (C<sub>5</sub>-C<sub>20</sub>) aryl, phenyl, phenyl, phena-1,4-diyl, naphthyl, naphtha-2,6-diyl, naphtha-2,7-diyl,  
30 (C<sub>6</sub>-C<sub>26</sub>) arylalkyl,  $-(CH_2)_i-\phi-$ ,  $-(CH_2)_i-\psi-$ ,  $-(CH_2)_i-NHR''-C(O)-\phi-$  and













$-(CH_2)_i-NHR''-C(O)-\psi-$ , where each  $i$  is independently an integer from 1 to 6,  $R''$  is hydrogen or  $(C_1-C_6)$  alkyl,  $\phi$  is  $(C_5-C_{20})$  aryl-diyl, phenyl-diyl or phena-1,4-diyl and  $\psi$  is naphthyl-diyl, naphtha-2,6-diyl or naphtha-2,7-diyl; and/or


$R_x$  is any electrophilic or nucleophilic group.

- 5 In yet another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ib) in which Y is selected from the group consisting of (Y-1), (Y-2), (Y-3), (Y-4), where substituent L is attached to  $R^3$ , or one of the following group of compounds, where  $R^9$  is as previously defined for structural formula (Y-1) and the dashed line at the nitrogen atom indicates the point of attachment of substituent L:

10 (Y-20b)	
15 (Y-21b)	
(Y-22b)	
20 (Y-23b)	

	(Y-24b)	
5	(Y-25b)	
	(Y-31b)	
10	(Y-34b)	
15	(Y-35b)	

(Y-36b)	
5 (Y-37b)	
(Y-39b)	
10 (Y-42b)	
15 (Y-43b)	



5

In still another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formula (Ib) in which:

Y is selected from the group consisting of (Y-1), (Y-2), (Y-3), (Y-4), (Y-20b), (Y-21b), (Y-22b), (Y-23b), (Y-24b), (Y-25b), (Y-31b), (Y-34b), (Y-35b), (Y-36b), (Y-39b), (Y-42b), (Y-43b), and (Y-46b); and

Z is selected from the group consisting of (C<sub>1</sub>-C<sub>12</sub>) alkyl, (C<sub>1</sub>-C<sub>12</sub>) alkanyl, (C<sub>5</sub>-C<sub>10</sub>) aryl, phenyl, naphthyl, naphth-1-yl, naphth-2-yl, pyridyl and purinyl, where in structures (Y-1) through (Y-4) L is attached to R<sup>3</sup>.


L is selected from the group consisting of (C<sub>1</sub>-C<sub>6</sub>) alkylidiyl, (C<sub>1</sub>-C<sub>6</sub>) alkano, (C<sub>5</sub>-C<sub>20</sub>) arylidiyl, phenylidiyl, phena-1,4-diyl, naphthylidiyl, naphtha-2,6-diyl, naphtha-2,7-diyl, (C<sub>6</sub>-C<sub>26</sub>) arylalkylidiyl  $-(CH_2)_i-\phi-$ ,  $-(CH_2)_i-\psi-$ ,  $-(CH_2)_i-NR''-C(O)-\phi-$  and  $-(CH_2)_i-NR''-C(O)-\psi-$ , where each *i* is independently an integer from 1 to 6, R'' is hydrogen or (C<sub>1</sub>-C<sub>6</sub>) alkyl,  $\phi$  is (C<sub>5</sub>-C<sub>20</sub>) arylidiyl, phenylidiyl or phena-1,4-diyl and  $\psi$  is naphthylidiyl, naphtha-2,6-diyl or naphtha-2,7-diyl, and where in structures (Y-1) through (Y-4), L is attached to R<sup>3</sup>.

R<sub>x</sub> is any electrophilic or nucleophilic group.

In still another preferred embodiment, the rhodamine dyes of the invention are compounds according to structural formulae (I), (Ia) and (Ib), including any of their respective previously-described preferred embodiments, in which E is a carboxyl or a salt thereof.

In the rhodamine dyes of the invention, aside from substituent E, the exact positions of the various substituents substituting the new bottom ring, *i.e.*, the Pyr<sup>+</sup> and  $-S-Z^1-L-R_x$  substituents illustrated in structural formula (Ia) and the Pyr<sup>+</sup> and  $-S-Z$  substituents illustrated in structural

formula (Ib) in the dyes of the present invention may be unknown. However, it has been confirmed by nuclear magnetic resonance (NMR) that the bottom ring contains one, two, or three  $\text{Pyr}^+$  substituents. Based upon the starting materials used to synthesize the rhodamine dyes of the invention and the nature of the synthesis reactions (discussed in more detail in Section 5.3.2, *infra*), it is believed that the various rhodamine dyes according to structural formula (I) have the regiochemistry depicted in structural formula (Ic) (illustrated with  $\text{Pyr}^+ =$  4-(dimethylamino)pyridinium and  $\text{E} = -\text{CO}_2\text{H}$ ):



It will be understood, however, that the Applicants do not intend to be bound by any particular structural representation, and that the invention is intended to encompass the rhodamine dyes that are obtained by the synthetic methods described herein, regardless of their specific regiochemistry.



In addition to the above-described structural isomerism about the new bottom ring, those of skill in the art will appreciate that many of the compounds encompassed by formulae (I), (Ia) and/or (Ib) as described herein, as well as many of the energy-transfer dye pairs and other conjugates described *infra*, may exhibit the phenomena of tautomerism, conformational isomerism, geometric isomerism and/or stereoisomerism. As the formulae drawings within this specification can represent only one of the possible tautomeric, conformational isomeric, enantiomeric or geometric isomeric forms, it should be understood that the invention encompasses any tautomeric, conformational isomeric, enantiomeric and/or geometric isomeric forms of the compounds having one or more of the utilities described herein.

Moreover, as the compounds of the invention may bear one or more positive charges or negative charges, depending upon their physical state, they may be in the form of a salts or may have counterions associated therewith. The identity(ies) of any associated counterions is typically dictated by the synthesis and/or isolation methods by which the compounds are obtained. Typical  
5 counterions include, but are not limited to, halides, acetate, trifluoroacetate, etc. and mixtures thereof. It will be understood that the identity(ies) of any associated counterions is not a critical feature of the invention, and that the invention encompasses any type of associated counterion. Moreover, as the compounds can exists in a variety of different forms, the invention is intended to encompass not only forms that are in association with counterions (*e.g.*, dry salts), but also forms  
10 that are not in association with counterions (*e.g.*, aqueous solutions).


### 5.3.2 Methods of Synthesis

The rhodamine dyes of the invention, and in particular rhodamine dyes according to structural formula (Ia) in which each  $\text{Pyr}^+$  is 4-(dimethylamino)pyridinium can be readily  
15 synthesized from corresponding tetrafluoro-rhodamine starting materials according to Scheme I, below. Although a specific regiochemistry is depicted in Scheme I, it will be understood that the structural representations are illustrative only. In Scheme I, Y, S, L,  $Z^1$  and  $R_x$  are as previously defined for structural formula (Ia). E is illustratively a carboxyl, but could be a sulfonate.

5




Scheme I



15

Referring to Scheme I, 4-(dimethyl)aminopyridine is added to a solution of tetrafluoro-rhodamine **100**. The reaction is followed by thin-layer chromatography (TLC). The compound will depend upon the identity of rhodamine-type parent xanthene ring Y. Once the reaction has gone to completion, thiol **108** is added to the mixture and the reaction monitored with TLC. Dye **110** is purified by reverse phase chromatography or other standard methods. Compounds in which  $\text{Pyr}^+$  is other than 4-(dimethylamino)pyridinium are prepared in an analogous manner from the appropriate aminopyridine starting material. Compounds having non-identical  $\text{Pyr}^+$  groups may be prepared using a mixture of the appropriate aminopyridines.

Compounds according to structural formula (**1b**) are synthesized in a similar manner from the appropriate tetrafluoro-rhodamine derivative **101** (illustrated below) using thiol Z-SH (**106**), where Z is as defined in structural formula (**1b**), to displace a  $\text{DMAP}^+$  group:

**101**

15

In derivative **101**, Y, L and  $\text{R}_x$  are as previously defined for structural formula (**1b**). When linking moiety  $-\text{L}-\text{R}_x$  is attached at a xanthene nitrogen, tetrafluoro-rhodamine **101** is prepared as illustrated in Scheme I. When linking moiety  $-\text{L}-\text{R}_x$  is attached at the xanthene C4 position, tetrafluoro-rhodamine **101** is prepared from a substituted monomer **118** ( $\text{R}^4 = \text{CH}_2\text{X}$  or  $\text{L}-\text{R}_x$ ). The full range of rhodamine dyes described herein can be synthesized by routine modification of these methods. Where necessary, any reactive functionalities on linker L, e.g. reactive functional group  $\text{R}_x$  can be protected using known groups and methods (see, e.g., Greene & Wuts, *Protective Groups in Organic Chemistry*, 1991, 2<sup>nd</sup> Edition, John Wiley & Sons, NY). Dyes in which  $\text{R}_x$  is a carboxylic acid or a salt thereof are particularly advantageous, as the carboxyl group does not require protection during synthesis.

The appropriate tetrafluoro-rhodamine starting materials can be produced in the usual



manner for the synthesis of rhodamines by condensing 1 mol of an appropriate 3-aminophenol derivative with 1 mol of tetrafluorophthalic anhydride **114** (Aldrich Chemical Co, St. Louis, MO) according to known techniques (*see, e.g.*, U.S. Patent No. 5,750,409; *Römpfs' Chemie Lexicon*, 8<sup>th</sup> Edition, pp. 3580). The appropriate 3-aminophenol starting materials are either commercially  
5 available or can be readily obtained using standard synthesis methods. A representative synthesis for preparing tetrafluororhodamine starting material **100** in which rhodamine-type parent xanthene ring Y is a compound according to structural formula (Y-1) (compound **120**) is outlined in Scheme II..




In Scheme II, the various R<sup>n</sup> substituents are as previously defined for structural formula (Y-1). According to Scheme II, 1 mol of 3-aminophenol derivative **112** is refluxed with 1 mol of tetrafluorophthalic anhydride **114** (Aldrich) to yield intermediate **116**. Intermediate **116** is refluxed with 3-aminophenol derivative **118** to yield tetrafluoro-rhodamine **120**. Depending upon the  
5 identity of rhodamine-type parent xanthene ring Y, the tetrafluoro-rhodamine **120** is isolated by reverse phase or normal phase column chromatography.

An analogous method for synthesizing tetrafluoro-rhodamine **100** in which rhodamine-type parent xanthene ring Y is a compound according to structural formula (Y-2) is illustrated in Scheme III:

10


Scheme III




In Scheme III, the various R<sup>n</sup> substituents are as previously defined for structural formula (Y-2). Dyes according to structural formula (Ia) in which the rhodamine-type parent xanthene ring is a compound according to structural formula (Y-3) or (Y-4) are prepared in an analogous manner from the appropriate starting materials.

- 5 Methods of preparing tetrafluoro-rhodamine derivative **101** in which Y is a compound according to structural formula (Y-1) or (Y-2) in which the linking moiety is attached to the xanthene nitrogen (*i.e.*, at position R<sup>3</sup>) are illustrated in Schemes IIb and IIIb, respectively:

Scheme IIb



Scheme IIIb



The various aminophenol starting materials are either commercially available or can be prepared using routine methods. Exemplary syntheses are provided in the Examples section. The full scope of the rhodamine compounds described herein can be readily synthesized by routine modification of any of these methods.

#### 5.4 Energy Transfer Dye Pairs


In another aspect, the present invention provides energy-transfer dye pairs incorporating the new rhodamine dyes of the invention. Generally, the energy-transfer dye pairs of the present invention comprise three main elements: (i) a donor dye (DD) which absorbs light at a first wavelength and emits excitation energy in response; (ii) an acceptor dye (AD) which is capable of absorbing the excitation energy emitted by the donor dye and emitting light at a second wavelength in response; and (iii) a linkage linking the donor dye to the acceptor dye, the linkage being effective to facilitate efficient energy-transfer between the donor and acceptor dyes. In the energy-transfer dye pairs of the invention, at least one of the donor or acceptor dyes, typically the acceptor dye, is a new rhodamine dye of the invention. The donor and acceptor dyes can be linked together in a variety of different configurations, depending upon the identities of the dyes. A thorough discussion of the various structures, synthesis and use of certain energy-transfer dye pairs

which may aid an understanding of the energy-transfer dye pairs of the invention is provided in U.S. Patent No. 5,800,996; U.S. Patent No. 5,863,727; and U.S. Patent No. 5,654,419, the disclosures of which are incorporated herein by reference in their entireties.

The energy-transfer dye pairs of the invention are typically obtained as illustrated in

# 5 Scheme V:

Scheme V



10

In Scheme V, a dye 140 which includes an optional linker L" and a complementary functional group F<sub>x</sub> is condensed with a rhodamine dye of the invention according to structural formula (Ia) or (Ib) to yield energy-transfer dye pairs according to structural formulae (IIIa) and (IIIb), respectfully, where for example three pyridinium rings are present..

During the condensation, reactive group R<sub>x</sub> and complementary functional group F<sub>x</sub> react to

form covalent linkage  $R^{41}$ . Thus, it will be recognized by those of skill in the art that reactive group  $R_x$  and complementary functional group  $F_x$  can each constitute respective members of the various pairs of complementary groups described in Section 5.3.1, such as various pairs of complementary electrophiles and nucleophiles. Preferably, one of  $R_x$  or  $F_x$  is an amine, thiol or hydroxyl group, most preferably an amine group, and the other one of  $R_x$  or  $F_x$  is a group capable of reacting with an amine, thiol or hydroxyl, most preferably a carboxylic acid or a salt, ester or activated ester thereof. Thus, a particularly preferred covalent linkage  $R^{41}$  is an amide of the formula  $-C(O)NR^{45}$ , where  $R^{45}$  is hydrogen or  $(C_1-C_6)$  alkyl.

In the energy-transfer dye pairs according to structural formulae **(IIa)** and **(IIb)**, DD/AD represents either a donor dye or an acceptor dye. Whether DD/AD constitutes a donor or acceptor will depend on the respective excitation and emission properties of DD/AD and Y. Typically, DD/AD is a dye belonging to the xanthene (including fluoresceine and rhodamine dyes), cyanine, phthalocyanine or squaraine classes of dyes. Alternatively, DD/AD can be a dye of the invention, preferably a dye according to structural formula **(Ia)** or **(Ib)**. The only requirement is that dye DD/AD either be capable of absorbing excitation energy emitted from Y or be capable of emitting excitation energy absorbable by Y. Preferably, in addition to complementary functional group  $F_x$ , DD/AD includes a linking moiety or linking group as previously described for conjugating the energy-transfer dye pairs of the invention to other compounds or substances.  $L''$  represents a bond or a linker analogous to the previously described linkers L that, when included in an energy-transfer dye pair according to structural formula **(IIa)** or **(IIb)**, facilitates efficient energy transfer between the acceptor and donor chromophores. The identity of  $L''$  and its point of attachment to DD/AD will depend, in part, upon the identity of DD/AD. Structures of these various classes of AD/DD dyes, as well as suitable linkers  $L''$  and points of attachment to these various classes of AD/DD dyes are described in U.S. Patent No. 5,863,727, the disclosure of which is incorporated herein by reference. When DD/AD is a cyanine dye,  $L''$  is preferably an amide, and is attached to the quaternary nitrogen. When DD/AD is a phthalocyanine dye,  $L''$  is preferably a sulfonamide, and is attached to the aromatic carbon skeleton. When DD/AD is a squaraine dye,  $L''$  is preferably an amide, and is attached to the quaternary nitrogen. When DD/AD is a rhodamine dye,  $L''$  is preferably an amide, and is attached to C4 or  $R^9$ . When DD/AD is a fluorescein dye,  $L''$  is preferably an amide, and is attached to C4 or  $R^9$ .

Most preferably, DD/AD is a donor dye which emits excitation energy absorbable by Y.



Preferred donor dyes are xanthene dyes, especially those which include a linking moiety or linking group on their bottom rings. Preferred amongst the xanthene dyes are the fluorescein dyes.

Fluorescein dyes suitable for use as donor dyes include, but are not limited to, the 4,7-


dichlorofluoresceins described in U.S. Patent No. 5,188,934, the extended fluoresceins described in

5 U.S. Patent No. 5,775,409, 5-carboxyfluorescein (5-FAM), 6-carboxyfluorescein (6-FAM), 5-(and 6)-carboxyfluorescein (5,6-FAM), NAN, Cl-FLAN and TET. Linker L" is typically attached to the xanthene C4 position of these preferred fluorescein donor dye. The actual choice of donor dye will depend upon the emission and excitation properties of DD/AD and Y, and will be apparent to those of skill in the art. In preferred fluorescein donor dyes that include a 5-, 6- or 5-(and 6)-carboxyl,


10 the carboxyl group (or an activated ester thereof) can be used to covalently conjugate the energy-transfer dye pairs of the invention to other compounds or substances.

The energy-transfer dyes of the invention are more fully described below with reference to various preferred embodiments. Referring to Scheme V, one group of preferred energy-transfer dye pairs according to structural formulae (IIa) and (IIb) are obtained when the compounds

15 according to structural formulae (Ia) and (Ib) are any of their previously-described preferred embodiments and compound 140 has the structure:



including any counter ions, wherein:



$R^{34}$ , when taken alone, is hydrogen, or when taken together with  $R^{35}$  is benzo;

$R^{35}$ , when taken alone, is hydrogen, fluoro, chloro, hydroxyl or carboxyl, or when taken together with  $R^{34}$  is benzo;

5  $R^{36}$  is hydrogen, fluoro or chloro;

$R^{37}$ , when taken alone, is hydrogen, fluoro, chloro, hydroxyl or carboxyl, or when taken together with  $R^{38}$  is benzo;

$R^{38}$ , when taken alone, is hydrogen, or when taken together with  $R^{37}$  is benzo;

$R^{39}$  is hydrogen or chloro;

10  $R^{40}$  is hydrogen or chloro; and

$R^{50}$  is carboxyl, or a salt, ester or activated ester thereof. Preferred compound **140** can be the pure 5-isomer, the pure 6-isomer or a mixture of 5-(and 6)-isomers. Particularly preferred compounds **140** are those in which  $R^{34}$ ,  $R^{35}$ ,  $R^{36}$ ,  $R^{37}$ ,  $R^{38}$ ,  $R^{39}$  and  $R^{40}$  are each hydrogen.

In another preferred embodiment, the energy-transfer dye pairs of the invention are

15 compounds according to structural formula (IIa) in which:

Y is a compound according to structural formula (Y-1), (Y-2), (Y-3), (Y-4), (Y-20a), (Y-21a), (Y-22a), (Y-23a), (Y-24a), (Y-25a), (Y-31a), (Y-34a), (Y-35a), (Y-36a), (Y-39a), (Y-41a), (Y-42a), (Y-43a), (Y-44a), (Y-45a), or (Y-46a);

20 DD/AD is a fluorescein dye having a 5- or 6-carboxyl group or a salt, ester or activated ester thereof, particularly a fluorescein dye selected from the group consisting of 5-carboxyfluorescein (5-FAM), 6-carboxyfluorescein (6-FAM), 5-(and 6)-carboxyfluorescein, NAN, CI-FLAN and TET;

L is a bond,  $(C_1-C_6)$  alkylidyl or  $(C_1-C_3)$  alkano, preferably a bond;

$Z^1$  is  $(C_1-C_6)$  alkylidyl,  $(C_5-C_{14})$  aryl or heteroaryl;

25  $R^{41}$  is  $-C(O)-R^{42}-$ , where  $R^{42}$  is O, S or NH and is bonded to L"; and/or

L" is  $-R^{43}-Z^3-C(O)-R^{44}-R^{45}-$ , wherein  $R^{43}$  is  $(C_1-C_6)$  alkylidyl, preferably  $(C_1-C_3)$  alkano, and is bonded to  $R^{42}$ ;  $Z^3$  is 5-6 membered cyclic alkenyldiyl or heteroalkenyldiyl,  $(C_5-C_{14})$  arylidyl or heteroaryldiyl;  $R^{44}$  is O, S or NH; and  $R^{45}$  is  $(C_1-C_6)$  alkylidyl, preferably  $(C_1-C_3)$  alkano, and is bonded to the xanthene C4 carbon of DD/AD.

Particularly preferred amongst the above-described energy-transfer dye pairs are those compounds in which:

L is a bond;

5  $Z^1$  and  $Z^3$  are each independently selected from the group consisting of phenyldiyl, phena-1,4-diyl, naphthyldiyl, naphtha-2,6-diyl and naphtha-3,6-diyl;

$R^{42}$  is NH;

$R^{43}$  is methano;

$R^{44}$  is NH; and/or

$R^{45}$  is methano.

10

In another preferred embodiment, the energy-transfer dye pairs of the invention are compounds according to structural formula (IIb) in which:

E is carboxyl or a salt thereof,

Y is a compound according to structural formula (Y-1), (Y-2), (Y-3) or (Y-4),

15 where L is attached at position  $R^3$ , or is a compound according to structural formula (Y-20b), (Y-21b), (Y-22b), (Y-23b), (Y-24b), (Y-25b), (Y-31b), (Y-34b), (Y-35b), (Y-36b), (Y-39b), (Y-41b), (Y-42b), (Y-43b), or (Y-46b);

DD/AD is a fluorescein dye having a 5- or 6-carboxyl group or a salt, an ester or an activated ester thereof, particularly a fluorescein dye selected from the group consisting of  
20 5-carboxyfluorescein (5-FAM), 6-carboxyfluorescein (6-FAM), 5-(and-6)-carboxyfluorescein, NAN, CI-FLAN and TET;

Z is  $(C_1-C_6)$  alkyl,  $(C_1-C_6)$  alkanyl,  $(C_5-C_{14})$  aryl or heteroaryl;

L is  $-CH_2-\phi-$  or  $-CH_2-\psi-$ , where  $\phi$  is phenyldiyl and  $\psi$  is naphthyldiyl and the methylene is attached to Y;

25

$R^{41}$  is  $-C(O)-R^{42}-$ , where  $R^{42}$  is O, S or NH; and/or

L" is  $-R^{43}-Z^3-C(O)-R^{44}-R^{45}-$ , wherein  $R^{43}$  is  $(C_1-C_6)$  alkyl, preferably  $(C_1-C_3)$  alkano, and is bonded to  $R^{42}$ ;  $Z^3$  is 5-6 membered cyclic alkenyldiyl or heteroalkenyldiyl,  $(C_5-C_{14})$  aryldiyl or heteroaryldiyl;  $R^{44}$  is O, S or NH; and  $R^{45}$  is  $(C_1-C_6)$  alkyl, preferably  $(C_1-C_3)$  alkano, and is bonded to the xanthene C4 carbon of DD/AD.

30

Particularly preferred amongst the above-described energy-transfer dye pairs are those compounds in which:

L is methano;

Z is selected from the group consisting of phenyl, naphthyl, naphth-1-yl and naphth-2-yl;

Z<sup>3</sup> is selected from the group consisting of phenyldiyl, phen-1,4-diyl, naphthyldiyl, naphth-2,6-diyl and naphth-3,6-diyl;

R<sup>42</sup> is NH;

R<sup>43</sup> is methano;

R<sup>44</sup> is NH; and/or

R<sup>45</sup> is methano.

10

#### **5.4.1 Synthesis of the Energy-transfer dye pairs**

Methods for synthesizing the various energy-transfer dyes of the invention are illustrated in Scheme V, *supra*. Conditions for carrying out the reactions are described in U.S. Patent No. 5,863,727, including Compound 140. Syntheses of exemplary energy-transfer dye pairs are described in the Examples section. All of the energy-transfer dye pairs of the invention can be obtained by routine modification of any of these methods.

15

#### **5.5 Conjugates Incorporating Dyes and Energy-transfer dye pairs**

In another aspect, the present invention comprises molecules and/or substances or conjugated with the rhodamine dyes and/or energy-transfer dye pairs of the invention. The conjugates can comprise virtually any molecule or substance to which the dyes or energy-transfer dye pairs of the invention can be conjugated, including by way of example and not limitation, peptides, polypeptides, proteins, saccharides, polysaccharides, nucleosides, nucleotides, polynucleotides, lipids, solid synthesis supports, organic and inorganic polymers, and combinations and assemblages thereof, such as chromosomes, nuclei, living cells (*e.g.*, bacteria or other microorganisms, mammalian cells, tissues, etc.), and the like. The dyes or energy-transfer dye pairs are conjugated with the reagent *via* a linking moiety by a variety of means, including hydrophobic attraction, ionic attraction, covalent attachment or with the aid of pairs of specific binding molecules, as previously described. Preferably, the dyes are conjugated *via* covalent attachment.

30

Conjugation typically results from mixing a dye or dye pair including an optional linking

moiety with and the molecule or substance to be conjugated in a suitable solvent in which both are soluble, using methods well-known in the art, followed by separation of the conjugate from any unconjugated starting materials or unwanted by-products. The dye or dye pair conjugate can be stored dry or in solution for later use.

5

### 5.5.1 Nucleoside/tide Conjugates

A preferred class of conjugates include nucleosides/tides and nucleoside/tide analogs that are labeled with the rhodamine dyes or energy-transfer dye pairs of the invention. Such labeled nucleosides/tides are particularly useful for labeling polynucleotides


10 formed by enzymatic synthesis, *e.g.*, labeled nucleotide triphosphates used in the context of template-directed primer extension, PCR amplification, Sanger-type polynucleotide sequencing, and/or nick-translation reactions. Labeled nucleoside/tides and/or nucleoside/tide analog conjugates are generally obtained by condensing a nucleoside/tide or nucleoside/tide analog (NUC) modified to contain a linking moiety ( $-L'-F_x$ ) with a rhodamine dye according to structural formula

15 **(Ia)** or **(Ib)** to yield a dye-labeled nucleoside/tide or nucleoside/tide analog according to structural formula **(IIIa)** or **(IIIb)**, respectively. Energy-transfer dye pair-labeled nucleosides/tides and/or nucleoside/tide analogs are obtained in an similar manner with energy-transfer dye compounds according to structural formulae **(IIa)** and **(IIb)** in which the DD/AD dye includes an optional linking moiety of the formula  $-L^3-R_x$ , where  $L^3$  is a bond or linker similar to previously-described




20 linker L (compounds **(IIa.1)** and **(IIb.1)**, respectively). These reactions are illustrated in Schemes VIa and VIb:

Scheme VIa

---






(IIIa)





(IIIb)

Scheme VIb



144



Referring to Schemes VIa and VIb, where for example three pyridinium rings are present and reactive functional group  $R_x$  on dyes or energy-transfer dye pairs according to structural formulae 5 (Ia), (Ib), (IIa.1) and/or (IIb.1) and complementary functional group  $F_x$  on compound 144 react to form covalent linkage  $R^{46}$ . Covalent linkage  $R^{46}$  is analogous to the previously-described covalent linkage  $R^{41}$ . Preferred embodiments of  $R^{46}$  are the same as the preferred embodiments previously described for  $R^{41}$ .


Complementary functional group  $F_x$  is attached to NUC *via* linker  $L'$ . Complementary 10 functional group  $F_x$  may be attached directly to NUC, in which case  $L'$  represents a bond, or it may be spaced away from NUC by one or more intervening atoms, in which case  $L'$  represents a linker. Any of the linkers  $L$  or  $L''$  previously described in connection with the rhodamine dye compounds *per se* and/or the energy-transfer dye pairs *per se* can be used for linker  $L'$ .

Complementary functional group  $F_x$  may be attached to NUC at a variety of different 15 positions, *e.g.*, the nucleobase, the sugar and/or the phosphate ester moiety. Nucleosides/tides and nucleoside/tide analogs that are appropriately modified at these various positions such that they can be conjugated with dyes and/or dye pairs according to the invention are known in the art. Preferably, complementary group  $F_x$  is attached to the nucleobase. When the nucleobase is a 7-deazapurine,  $L'$  is usually attached to the C7 position of the nucleobase. When the nucleobase is a 20 pyrimidine,  $L'$  is usually attached to the C-5 position of the nucleobase. When the nucleobase is a purine,  $L'$  is usually attached to the C-8 position of the nucleobase. Linkers  $L'$  useful for covalently conjugating the dyes of the invention to the nucleobase of NUC are described in U.S. Patent No. 5,821,356, U.S. Patent No. 5,770,716 and U.S. application Serial No. 08/833,854 filed April 10, 1997, the disclosures of which are incorporated herein by reference.

25 Preferred linkers  $L'$  for covalently conjugating the dyes of the invention to the nucleobase of NUC include ( $C_2$ - $C_{20}$ ) alkylenos, heteroalkyldiyls and heteroalkylenos, especially ( $C_2$ - $C_{20}$ ) alkynos, ( $C_2$ - $C_{20}$ ) alkenos, heteroalkynos and heteroalkenos. A particularly preferred linker  $L'$  is  $-C\equiv C-CH_2-$ , where the terminal *sp* carbon is covalently attached to the nucleobase of NUC and the terminal methylene (*sp*<sup>3</sup>) carbon is covalently attached to  $R^{46}$  in the compounds of structural 30 formulae (IIIa), (IIIb), (IVa) and (IVb), or to  $F_x$  of compound 144.

Additional preferred linkers  $L'$  for covalently conjugating the rhodamine dyes or energy-transfer dye pairs of the invention to the nucleobase of NUC include propargylethoxy groups




according to structural formula  $-C\equiv C-CH_2-O-CH_2-CH_2-NR^{47}-R^{48}-$ , wherein  $R^{47}$  is hydrogen or  $(C_1-C_6)$  alkyl and  $R^{48}$  is selected from the group consisting of  $-C(O)-(CH_2)_r-$ ,  $-C(O)-CHR^{49}-$ ,  $-C(O)-C\equiv C-CH_2-$  and  $-C(O)-\phi-(CH_2)_r-$ , where each  $r$  is independently an integer from 1 to 5 and  $\phi$  represents a  $C_6$  arylidyl or heteroarylidyl, preferably phena-1,4-diyl (  ) and  $R^{49}$  is

- 5 hydrogen,  $(C_1-C_6)$  alkyl or an amino acid side chain (including side chains of both gene-encoded and non-encoded amino acids). With these linkers  $L'$ , the terminal  $sp$  carbon is attached to the nucleobase of NUC and the other terminal group is attached to  $R^{46}$  in the compounds of structural formulae (IIIa), (IIIb), (IVa) and (IVb), or to  $F_x$  of compound 144.

- In a preferred embodiment, the labeled nucleosides/tide and/or nucleoside/tide analogs  
10 according to structural formulae (IIIa), (IIIb), (IVa) and (IVb) are labeled enzymatically-incorporatable nucleotides, labeled enzymatically extendable nucleotides or labeled terminators.

In another preferred embodiment, the labeled nucleosides/tides and nucleoside/tide analogs are those obtained from Schemes (VIa) and (VIb) in which the compounds according to structural formulae (Ia), (Ib) are any of their respective previously-defined preferred embodiments,

- 15 compounds (IIa.1) and (IIb.1) are any of the previously defined preferred embodiments of compounds (IIa) and (IIb), respectively, and compound 144 is a compound according to structural formula (V):



20

wherein:

B is a nucleobase;


$F_x$  is a complementary functional group as previously described;

$L'$  is one of the preferred linkers described above;

25

$R_{70}$  and  $R_{71}$ , when taken alone, are each independently selected from the group consisting of hydrogen, hydroxyl and a moiety which blocks polymerase-mediated template-directed nucleic acid synthesis, or when taken together form a bond such that the illustrated sugar is 2',3'-didehydroribose; and

$R_{72}$  is selected from the group consisting of hydroxyl, a phosphate ester having



analog. Typically,  $F_x$  is an amino group of the formula  $\text{---NHR}^{51}$ , where  $R^{51}$  is hydrogen or  $(C_1\text{--}C_6)\text{alkyl}$ , but can be any nucleophilic or electrophilic groups.

10 In a preferred embodiment of structural formula (V), B is a normal nucleobase or a common analog thereof, a 7-deazapurine, 8-aza,7-deazapurine, a purine or a pyrimidine. In a particularly preferred embodiment, B is a nucleobase selected from the group consisting of 7-deaza-adenine, cytosine, 7-deaza-guanine, thymine and uracil. When the preferred nucleobase B is a purine or a 7-deaza-purine, the pentose moiety is attached to the  $N^9$ -position of the nucleobase, and when the preferred B is a pyrimidine, the pentose moiety is attached to the  $N^1$ -position of the nucleobase. Linker L' is attached to nucleobase B as previously described.

In structural formula (V), when both  $R_{70}$  and  $R_{71}$  are hydroxyl, the resultant compounds produced in Schemes VIa and VIb are labeled ribonucleoside/tides. When  $R_{70}$  is hydrogen and  $R_{71}$  is hydroxyl, the resultant compounds are labeled 2'-deoxyribonucleoside/tides. When  $R_{70}$  and  $R_{71}$  are each hydrogen, the resultant compounds are 2',3'-dideoxyribonucleoside/tides. Labeled 2',3'-dideoxyribonucleoside-5'-triphosphates (ddNTPs) find particular use as terminators in Sanger-type DNA sequencing methods utilizing fluorescent detection. Labeled 2'-deoxyribonucleoside-5'-triphosphates (dNTPs) find particular use as means for labeling DNA polymerase extension products, *e.g.*, in the polymerase chain reaction or nick-translation. Labeled ribonucleoside-5'-triphosphates (NTPs) find particular use as means for labeling RNA polymerase extension products.

Referring to Schemes (VIa) and (VIb), *supra*, the synthesis of alkynylamino-derivatized

compounds **144** useful for conjugating the dyes of the invention to nucleosides/tides and/or nucleoside/tide analogs is taught in EP 87305844.0 and Hobbs *et al.*, 1989, J. Org. Chem. 54:3420. The corresponding nucleoside mono-, di- and triphosphates are obtained by standard techniques (*see, e.g.*, the methods described in U.S. Patent No. 5,821,356, U.S. Patent No. 5,770,716 and U.S. application Serial No. 08/833,854 filed April 10, 1997, discussed *supra*). Methods for synthesizing compound **144** in modified with propargylethoxyamido linkers L' can also be found in these patents and application.

Energy-transfer dye pairs can be conjugated to a nucleotide 5'-triphosphate **144** by linking through a nucleobase amino group to: (i) an activated ester of a energy-transfer dye pair, e.g. **230** NHS ester, or (ii) stepwise coupling to one dye **140**, e.g. 4'-protected aminomethyl fluorescein, then coupling the unprotected 4'-aminomethyl to the second dye of the pair, e.g. **196** NHS ester.

Additional synthesis procedures suitable for use in synthesizing compounds according to structural formulae (**IIIa**), (**IIIb**), (**IVa**) and (**IVb**) are described, for example, in Gibson *et al.*, 1987, Nucl. Acids Res. 15:6455-6467; Gebeyehu *et al.*, 1987, Nucl. Acids Res. 15:4513-4535; Haralambidis *et al.*, 1987, Nucl. Acids Res. 15:4856-4876; Nelson *et al.*, 1986, Nucleosides and Nucleotides. 5(3):233-241; Bergstrom *et al.*, 1989, J. Am. Chem. Soc. 111:374-375; U.S. Patent No. 4,855,225, U.S. Patent No. 5,231,191 and U.S. Patent No. 5,449,767, the disclosures of which are incorporated herein by reference. Any of these methods can be routinely adapted or modified as necessary to synthesize the full range of labeled nucleosides/tides and nucleoside/tide analogs described herein.

#### 5.5.1.1 Polynucleotide Conjugates

Yet another preferred class of conjugates of the present invention comprise polynucleotides and/or polynucleotide analogs labeled with the rhodamine dyes or energy-transfer dye pairs of the invention. Such labeled polynucleotides and/or analogs are useful in a number of important contexts, including as DNA sequencing primers, PCR primers, oligonucleotide hybridization probes, oligonucleotide ligation probes, and the like.

In one preferred embodiment, the labeled polynucleotides or polynucleotide analogs of the present invention include multiple dyes located such that fluorescence energy-transfer takes place between a donor dye and an acceptor dye. Such multi-dye energy-transfer polynucleotides find application as spectrally-tunable sequencing primers as described, for example, in Ju *et al.*, 1995, Proc. Natl. Acad. Sci. USA 92:4347-4351, and as hybridization probes as described, for example,

in Lee *et al.*, 1993, Nucl. Acids Res. 21:3761-3766.

Labeled polynucleotides and/or polynucleotide analogs are typically synthesized enzymatically, *e.g.*, using a DNA/RNA polymerase or ligase (*see, e.g.*, Stryer, 1981, *Biochemistry*, Chapter 24, W.H. Freeman and Company) and a labeled enzymatically-incorporatable nucleotide, 5 as previously described. Alternatively, the labels may be introduced subsequent to synthesis *via* conventional conjugation reactions, as discussed more thoroughly below.

Generally, if the labeled polynucleotide is made using enzymatic synthesis, the following procedure may be used. A target DNA polynucleotide is denatured and an oligonucleotide primer is annealed to the target DNA. A mixture of 2'-deoxyribonucleoside-5'-triphosphates capable of 10 supporting template-directed enzymatic extension of the primed target (*e.g.*, a mixture including dGTP, dATP, dCTP, and dTTP) is added to the primed target. At least a fraction of the deoxynucleotides are labeled with a rhodamine dye or energy-transfer dye pair of the invention as described above. Next, a polymerase enzyme is added to the mixture under conditions where the polymerase enzyme is active. A labeled polynucleotide is formed by the incorporation of the 15 labeled deoxynucleotides during polymerase-mediated strand synthesis. In an alternative enzymatic synthesis method, two primers are used instead of one: one complementary to the plus (+) strand of the target and another complementary to the minus (-) strand of the target, the polymerase is a thermostable polymerase and the reaction temperature is cycled between a denaturation temperature and an extension temperature, thereby exponentially synthesizing a 20 labeled complement to the target sequence by PCR (*see, e.g.*, *PCR Protocols*, 1990, Innis *et al.* Eds., Academic Press).

Alternatively, the labeled polynucleotide or polynucleotide analog is obtained *via* post-synthesis conjugation. According to this embodiment, a polynucleotide or polynucleotide analog which includes a complementary functional group  $F_x$ , typically an amino group, is condensed with 25 a rhodamine dye according to structural formula (IIa) or (IIb) or an energy-transfer dye pair according to structural formula (IIIa) or (IIIb) under conditions wherein  $R_x$  and  $F_x$  react to form a covalent linkage. The labeled polynucleotides and/or polynucleotide analogs are isolated using conventional means, such as alcohol precipitation, gel electrophoresis, column chromatography, etc.

30 A variety of reagents for introducing amino groups, or other complementary functional groups such as thiols into enzymatically or chemically synthesized polynucleotides and polynucleotide analogs are known in the art, as are appropriate condensation conditions. For

example, methods for labeling polynucleotides at their 5'-terminus are described in *Oligonucleotides and Analogs*, 1991. Eckstein, Ed., Chapter 8, IRL Press; Orgel *et al.*, 1983, Nucl. Acids Res. 11(18):6513; and U.S Patent No. 5,118,800. Methods for labeling polynucleotides at the phosphate ester backbone are described in *Oligonucleotides and Analogs*, 1991. Eckstein, Ed., Chapter 9, IRL Press. Methods for labeling polynucleotides at their 3'-terminus are described in Nelson *et al.*, 1992, Nucl. Acids Res. 20(23):6253-6259; U.S. Patent No. 5,401,837; and U.S. Patent No. 5,141,813. For a review of labeling procedures, the reader is referred to Haugland, *In: Excited States of Biopolymers*, 1983, Steiner. Ed., Plenum Press, NY. All of these disclosures are incorporated herein by reference.

## 5.6 Methods Utilizing the Dyes and Reagents of the Invention

The rhodamine dyes, energy-transfer dye pairs and conjugates incorporating the dyes and energy-transfer dye pairs of the present invention are well suited to any method utilizing fluorescent detection, particularly aqueous applications and methods requiring the simultaneous detection of multiple spatially-overlapping analytes. The various dyes and conjugates of the invention are particularly well suited for identifying classes of polynucleotides that have been subjected to a biochemical separation procedure, such as electrophoresis, or that have been distributed among locations in a spatially-addressable hybridization array.

The various dyes of the invention can be conjugated to peptides, proteins, antibodies, and antigens. Dye-antibody conjugates are useful for sandwich-type immunosorbent assays. Especially preferred are bead-based assays for detection of peptides, cells, and other cellular components where the dyes of the invention are conjugated to antibodies.

In a preferred category of methods referred to herein as "fragment analysis" or "genetic analysis" methods, labeled polynucleotide fragments are generated through template-directed enzymatic synthesis using labeled primers or nucleotides, *e.g.*, by ligation or polymerase-directed primer extension; the fragments are subjected to a size-dependent separation process, *e.g.*, electrophoresis or chromatography; and, the separated fragments are detected subsequent to the separation, *e.g.*, by laser-induced fluorescence. In a particularly preferred embodiment, multiple classes of polynucleotides are separated simultaneously and the different classes are distinguished by spectrally resolvable labels.

One such fragment analysis method known as amplified fragment length polymorphism detection (AmpFLP) is based on amplified fragment length polymorphisms, *i.e.*, restriction

fragment length polymorphisms that are amplified by PCR. These amplified fragments of varying size serve as linked markers for following mutant genes through families. The closer the amplified fragment is to the mutant gene on the chromosome, the higher the linkage correlation. Because genes for many genetic disorders have not been identified, these linkage markers serve to help evaluate disease risk or paternity. In the AmpFLPs technique, the polynucleotides may be labeled by using a labeled polynucleotide PCR primer, or by utilizing labeled nucleotide triphosphates in the PCR.

In another such fragment analysis method known as nick translation, a reaction is used to replace unlabeled nucleotides in a double-stranded (ds) DNA molecule with labeled nucleotides.

Free 3'-hydroxyl groups are created within the dsDNA by "nicks" caused by treatment with deoxyribonuclease I (DNAase I). DNA polymerase I then catalyzes the addition of a labeled nucleotide to the 3'-hydroxyl terminus of the nick. At the same time, the 5' to 3'-exonuclease activity of this enzyme eliminates the nucleotide at the 5'-phosphoryl terminus of the nick. A new nucleotide with a free 3'-OH group is incorporated at the position of the original excised nucleotide, and the nick is shifted along by one nucleotide in the 3' direction. This 3' shift will result in the sequential addition of new labeled nucleotides into the dsDNA. The nick-translated polynucleotide is then analyzed, for example, by using a separation process such as electrophoresis.

Another exemplary fragment analysis method is based on variable numbers of tandem repeats, or VNTRs. VNTRs are regions of double-stranded DNA that contain multiple adjacent copies of a particular sequence, with the number of repeating units being variable. Examples of VNTR loci are pYNZ22, pMCT118, and Apo B. A subset of VNTR methods are those methods based on the detection of microsatellite repeats, or short tandem repeats (STRs), *i.e.*, tandem repeats of DNA characterized by a short (2-4 bases) repeated sequence. One of the most abundant interspersed repetitive DNA families in humans is the (dC-dA)<sub>n</sub>-(dG-dT)<sub>n</sub> dinucleotide repeat family (also called the (CA)<sub>n</sub> dinucleotide repeat family). There are thought to be as many as 50,000 to 100,000 (CA)<sub>n</sub> repeat regions in the human genome, typically with 15-30 repeats per block. Many of these repeat regions are polymorphic in length and can therefore serve as useful genetic markers. Preferably, in VNTR or STR methods, label is introduced into the polynucleotide fragments using a labeled PCR primer.

In a particularly preferred fragment analysis method, classes identified in accordance with the invention are defined in terms of terminal nucleotides so that a correspondence is established between the four possible terminal bases and the members of a set of spectrally resolvable dyes.

Such sets are readily assembled from the dyes and energy-transfer dye pairs of the invention by measuring emission and absorption bandwidths with commercially available spectrophotometers. More preferably, the classes arise in the context of the chemical or chain termination methods of DNA sequencing, and most preferably the classes arise in the context of the chain termination  
5 methods, *i.e.*, dideoxy DNA sequencing, or Sanger-type sequencing.

Sanger-type sequencing involves the synthesis of a DNA strand by a DNA polymerase *in vitro* using a single-stranded or double-stranded DNA template whose sequence is to be determined. Synthesis is initiated at a defined site based on where an oligonucleotide primer anneals to the template. The synthesis reaction is terminated by incorporation of a terminator that  
10 will not support continued DNA elongation. When proper proportions of dNTPs and a single terminator complementary to A, G, C or T are used, enzyme-catalyzed primer extension will be terminated in a fraction of the extension products at each site where the terminator is incorporated. If labeled primers or labeled terminators are used for each reaction, the sequence information can be detected by fluorescence after separation of the resultant primer extension products by  
15 high-resolution electrophoresis. In the chain termination method, dyes or energy-transfer dye pairs of the invention can be attached to either the sequencing primers or terminators. The dyes or energy-transfer dye pairs can be linked to a complementary functionality on the 5'-end of the primer, *e.g.* following the teaching in Fung *et al.*, U.S. Patent No. 4,757,141; on the base of a primer; or on the base of a terminator, *e.g.* via the alkynylamino or other linking groups described  
20 above. Concentration ranges for the various enzymes, primers, dNTPs and labeled terminators are those commonly employed in the art.

In each of the above fragment analysis methods, labeled extension products are preferably separated by electrophoretic procedures, *e.g.* *Gel Electrophoresis of Nucleic Acids: A Practical Approach*, 1981, Rickwood and Hames, Eds., IRL Press Limited, London; Osterman, 1984,  
25 *Methods of Protein and Nucleic Acid Research*, Vol. 1 Springer-Verlag, Berlin; or U.S. Patent Nos. 5,374,527, 5,624,800 and/or 5,552,028. Preferably, the type of electrophoretic matrix is crosslinked or uncrosslinked polyacrylamide having a concentration (weight to volume) of between about 2-20 weight percent. More preferably, the polyacrylamide concentration is between about 4-8 percent. Preferably, in the context of DNA sequencing, the electrophoresis matrix includes a  
30 denaturing agent, *e.g.*, urea, formamide, and the like. Detailed procedures for constructing such matrices are given by Maniatis *et al.*, 1980, "Fractionation of Low Molecular Weight DNA and RNA in Polyacrylamide Gels Containing 98% Formamide or 7 M Urea," *Methods in Enzymology*

65:299-305; Maniatis *et al.*, 1975, "Chain Length Determination of Small Double- and Single-Stranded DNA Molecules by Polyacrylamide Gel Electrophoresis," *Biochemistry* 14:3787-3794; Maniatis *et al.*, 1982, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, pgs. 179-185; and *ABI PRISM™ 377 DNA Sequencer User's*  
5 *Manual, Rev. A*, January 1995, Chapter 2 (p/n 903433, The Perkin-Elmer Corporation, Foster City, CA). The optimal electrophoresis conditions, *e.g.*, polymer concentration, pH, temperature, and concentration of denaturing agent, employed in a particular separation depend on many factors, including among others, the size range of the nucleic acids to be separated, their base compositions, whether they are single stranded or double stranded, and the nature of the classes for which  
10 information is sought by electrophoresis. Accordingly, application of the invention may require standard preliminary testing to optimize conditions for particular separations.

Subsequent to electrophoretic separation, the labeled extension products are detected by measuring the fluorescence emission from the labels. To perform such detection, the labeled products are illuminated by standard means, *e.g.* high intensity mercury vapor lamps, lasers, or the  
15 like. The illumination wavelength will depend upon the spectral properties of the particular label. Preferably, the illumination means is a laser having an illumination beam at a wavelength between 400 and 700 nm. More preferably, the illumination means is a laser light generated by an argon ion laser, particularly the 488 and 514 nm emission lines of an argon ion laser, or the 532 emission line of a neodymium solid-state YAG laser or the 633 nm emission line of a helium-neon laser. Several  
20 argon ion lasers are available commercially which lase simultaneously at these lines, *e.g.* Cyonics, Ltd. (Sunnyvale, Calif.) Model 2001, or the like. The fluorescence is then detected by a light-sensitive detector, *e.g.*, a photomultiplier tube, a charged coupled device, or the like. Suitable exemplary electrophoresis detection systems are described elsewhere, *e.g.*, U.S. Patent Nos. 5,543,026; 5,274,240; 4,879,012; 5,091,652 and 4,811,218.

25 In preferred embodiments, the primer is unlabeled and the sequencing reaction includes, in addition to the polymerase and mixture of dNTPs, a mixture of four different terminators, one complementary to A, one complementary to G, one complementary to C and one complementary to T. Each of the different terminators is labeled with a different, spectrally resolvable dye or energy-transfer dye pair. One of the terminators is labeled with a rhodamine dye or energy-transfer  
30 dye pair of the invention. As each of the labeled terminators fluoresces at a different wavelength, following separation based on size, the identity of the 3'-terminal nucleotide of each extension product is identified by the wavelength (or color) of the label. In particularly preferred



embodiments, each of the different spectrally resolvable labels can be excited using a single light source. A set of such preferred labeled terminators is provided in the Examples section. Other sets will depend upon the excitation and emission spectral properties of the various labels are readily obtained by routine methods.

5

The invention having been described, the following examples are provide to illustrate, and not limit, the invention.

## 6. EXAMPLE: SYNTHESIS OF RHODAMINE DYE 190


10 Rhodamine dye 190 was synthesized as illustrated in Schemes I and II, *supra*, from the appropriate aminophenol starting materials.

15


20

25


30



### 6.1 Synthesis of 202




10 Amino-phenol compound **200** was synthesized as described in U.S. Patent No. 5,750,409 by alkylation of **192** with ethyl iodide and sodium bicarbonate in acetonitrile followed by demethylation with boron tribromide in dichloromethane.



15 202

A solution of aminophenol **200** (4.8 g, 22 mmol) and tetrafluorophthalic anhydride **114** (4.9 g, 22 mmol; Aldrich) was refluxed in toluene (44 ml) for 3 hr. The solution was cooled to rt and the precipitate collected to yield tetrafluoro ketone **202** (6.8 g, 70%).

## 6.2 Synthesis of 204




204

5 A suspension of the tert-butyl ester amine **194**, prepared according to U.S. Patent No. 5,688,808, (5.46 g, 20 mmol), methyl 4-(bromomethyl)benzoate (8.3 g, 36.2 mmol), sodium iodide (3.0 g, 20 mmol) and potassium carbonate (2.8 g, 20 mmol) was refluxed in acetonitrile (20 ml) for 1 hr. TLC (hexane/ethyl acetate 1/1) showed the reaction was complete. Ether was added to the cooled mixture and the solution was decanted from the solids. The solution was washed with

10 water (2 portions of 50 ml), brine (50 ml) and dried over magnesium sulfate. The crude product was purified by chromatography on silica gel (hexane/ethyl acetate 19:1) to give white crystals of the alkylated, tert-butyl ester product (4.3 g, 10 mmol, 50%). The tert-butyl ester was dissolved in a solution of lithium hydroxide monohydrate (1.7 g, 40 mmol) in water (10 ml) and methanol (50 ml), refluxed for 2 hr and then evaporated to dryness under vacuum. The residue was extracted

15 with diethylether (100 ml) and washed with water and brine. The organic layer was dried over magnesium sulfate, filtered and evaporated to give **204** as a tan foam.  $^1\text{H}$  NMR (**204**,  $\text{CD}_2\text{Cl}_2$ )  $\delta$  8.03, 2H, d; 7.45, 2H, d; 6.96, 1H, d; 6.04, 1H, d; 5.65, 1H, s; 4.55, 1H, m; 1.38, 3H, s; 1.25, 6H, s.

### 6.3 Synthesis of 206

**206**

5 Phosphorous oxytrichloride (2.8 ml, 30 mmol) was added to a suspension of tetrafluoro ketone **202** (4.4 g, 10 mmol) in chloroform (100 ml). The suspension was stirred at rt for 30 min., aminophenol **204** (3.2 g, 10 mmol) was added and the mixture was refluxed for 3 hr. The solution was cooled to rt and the reaction quenched with water (1 ml). The solvent was evaporated and dye **206** was purified by normal phase chromatography (DCM/MeOH/HOAc, 90:10:1). Dye **206** was  
10 further purified by C18 reverse phase chromatography (MeOH / 0.1 M TEAA, 4:1) to afford a dark green solid (0.65 g, 9%, Abs. max 610 nm, Em. max 632 nm, H<sub>2</sub>O).

#### 6.4 Synthesis of Dye 190

Dimethylaminopyridine (85 mg) was added to tetrafluoro dye **206** (30 mg) in 0.7 ml of  
15 dimethylformamide. After 20 hours at room temperature, TLC analysis (CH<sub>2</sub>Cl<sub>2</sub>:CH<sub>3</sub>OH:CH<sub>3</sub>CO<sub>2</sub>H 80:20:16) showed the complete disappearance of starting **206** and the appearance of a new, lower R<sub>f</sub> spot, the tetradimethylaminopyridinium adduct. Thiophenol (0.75 ml) was added, and after another 8 hours, TLC showed partial conversion to a higher R<sub>f</sub> spot. After concentration under vacuum, **190** was purified by reverse-phase HPLC (C18) by a gradient of  
20 30-50% acetonitrile in 0.1M TEAA. The fractions containing **190** were combined and evaporated to give a blue oil which was precipitated in either to provide pure dye **190** as a blue solid (30 mg, Abs. Max 628 nm, Em. max 650 nm, H<sub>2</sub>O).

The succinimidyl (NHS) ester of **190** was prepared from a solution of **190** (5 mg) in 100 µl


dimethylformamide (DMF). Succinimidyl tetramethyluronium tetrafluoroborate (20mg) and diisopropylamine (10  $\mu$ l) were added. After 1 hour at room temperature, TLC analysis (C2-reverse phase plates, CH<sub>3</sub>OH / 0.1M TEAA 1:1) showed the disappearance of starting **190** and a new, lower R<sub>f</sub> spot. The NHS ester of **190** was isolated by precipitation as a blue solid.

5

## 7. EXAMPLE: SYNTHESIS OF RHODAMINE DYE 196


Rhodamine dye **196** was synthesized in a manner analogous to dye **190** from the corresponding tetrafluoro dye **220**.

10



196

### 7.1 Synthesis of Tetrafluoro 220

**220**

5

Phosphorous oxytrichloride (2.8 ml, 30 mmol) was added to a suspension of tetrafluoro ketone **202** (4.4 g, 10 mmol) in chloroform (100 ml). The suspension was stirred at rt for 30 min., aminophenol **200** (3.2 g, 10 mmol) was added and the mixture was refluxed for 3 hr. The solution was cooled to rt and the reaction quenched with water (1 ml). The solvent was evaporated and dye **220** was purified by normal phase chromatography (DCM/MeOH/HOAc, 90:10:1). Dye **220** was further purified by C18 reverse phase chromatography (CH<sub>3</sub>OH : 0.1M TEAA, 4:1) to afford a dark green solid (0.65 g, 9%, Abs. max 613 nm, Em. max 643 nm, H<sub>2</sub>O).

## 7.2 Synthesis of Rhodamine Dye 196



To a solution of **220** (100 mg) in dimethylformamide (1.5 mL) was added 4-(dimethylamino)pyridine (150 mg). Thin-layer chromatography (TLC) on silica gel eluting with 80:20:16 dichloromethane:methanol:acetic acid could distinguish tetrafluoro **220** (R<sub>f</sub> = 1) from the tetra-dimethylaminopyridinium adduct (R<sub>f</sub> = 0). After 40 hr, analysis by thin-layer chromatography showed complete conversion to the SymJAZ adduct.

To this solution, 4-carboxythiobenzene (50 mg) was added. After 10 min, TLC analysis showed conversion to rhodamine dye **196** (R<sub>f</sub> = 0.1). Purification of **196** was accomplished on C18 silica gel with stepwise elution with 20-70% methanol vs. 0.1 M triethylammonium acetate. The dye **196** eluted between 30% and 50% methanol. The solvent was evaporated and the residual blue oil was precipitated with ether to provide the dye **196** as a blue solid (85 mg, Abs. max 636


nm, Em. max 661 nm, H<sub>2</sub>O).

### 7.3 Synthesis of Tetrafluoro 198

5



Amino-phenol compound **228** was synthesized as described in U.S. Patent No. 5,750,409 and cyclized with tetrafluorophthalic anhydride **114** as in Example 6.1 *supra* to give **238** (lambda max 384 nm). The mixture of 2.6 mmole **238** and 2.6 mmole **228** in 1.2 gm phosphorus oxytrichloride and 10 ml acetonitrile was refluxed for 12 hours. The solvent was evaporated under vacuum, dissolved in several milliliters of dichloromethane and chromatographed on silica gel, eluting with dichloromethane and methanol. The product fractions were combined, evaporated and further purified by reverse-phase preparative HPLC and precipitation in 1% HCl to give 35 mg **198** as a red solid. <sup>1</sup>H NMR (**198**, CD<sub>3</sub>OD) δ 6.82, 2H, s; 6.59, 2H, s; 5.63, 2H, s; 1.9, 6H, br s; 1.4, 12H, br s.




198

5

#### 7.4 Synthesis of Tetrafluoro 240


Tetrafluoro **240** was synthesized by the method and corresponding methylated intermediates of Example 7.3. Mass spectroscopy **240**: Exact Mass = 662.22 (Molecular Wt. = 662.67) Ex. max 624 nm, Em. max 644 nm (8M urea).



240

#### 15 8. EXAMPLE: SYNTHESIS OF RHODAMINE DYE 232






5

Rhodamine dye **232** was synthesized as illustrated in Schemes (I) and (II), *supra*, from the appropriate aminophenol starting materials.

#### 8.1 Synthesis of tricyclic amine-phenol **208**


10



A suspension of the tert-butyl ester amine **194**, prepared according to U.S. Patent No. 5,688,808, (12.8 g, 47 mmol), 1-bromo-3-chloropropane (29.3 g, 187 mmol), sodium iodide (56.4


g, 376 mmol) and sodium bicarbonate (7.9 g, 94 mmol) was refluxed in acetonitrile (150 ml) for 18 hr. The mixture was cooled to room temperature, filtered by suction filtration and evaporated to leave a residue. The filter cake was washed with hexane (300 ml) and the filtrate was combined with the residue and washed with water (2 portions of 50 ml), brine (50 ml) and dried over magnesium sulfate. The crude cyclized product was purified by chromatography on silica gel (hexane/ethyl acetate 20:1) to leave the intermediate tricyclic pivalate ester as a pale yellow oil (9.5 g, 30 mmol, 64%). The cyclized ester was dissolved in a solution of lithium hydroxide monohydrate (2.6 g, 60 mmol) in water (15 ml) and methanol (120 ml). The solution was stirred at room temperature for 1 hr and then evaporated to dryness under vacuum. The residue was treated with 1 M HCl (30 ml) and extracted with diethylether (3 portions of 100 ml). The combined extracts were washed with 200 mM pH 7 phosphate buffer (50 ml), dried over magnesium sulfate, filtered and evaporated to give tricyclic amine-phenol **208** as a brown solid, used crude in subsequent steps.

## 8.2 Synthesis of Tetrafluoro Ketone **210**



A solution of aminophenol **208** (0.67 g, 2.9 mmol) and tetrafluorophthalic anhydride (0.64 g, 2.9 mmol) was refluxed in toluene (10 ml) for 3 hr. The solution was cooled to rt and the precipitate collected, yielding tetrafluoro ketone **210** (0.92 g, 71 %).

## 8.3 Synthesis of Tetrafluoro **212**




Phosphorous oxytrichloride (0.56 ml, 6 mmol) was added to a solution of F4 RAZ ketone **210** (0.91 g, 2 mmol) in chloroform (20 ml). The solution was stirred for 15 min., aminophenol **208** (0.46 g, 2 mmol) was added and the mixture was refluxed for 3 hr. The solution was cooled to rt and the reaction quenched with water (0.5 ml). Tetrafluoro dye **212** was purified by normal phase chromatography (DCM/MeOH, 20:1) and further purified by C18 reverse phase chromatography (MeOH / 0.1 M TEAA, 9:1) to afford **212** as a metallic green solid (0.39 g, 30%, Abs. max 630 nm, Em. max 655 nm, 8M urea).

#### 8.4 Synthesis of Rhodamine Dye 232

Rhodamine dye **232** was synthesized from tetrafluoro dye **212** with 4-(dimethylamino)pyridine and 4-carboxythiobenzene, as described in Section 7.2, *supra*, and purified by C-8 reverse phase HPLC, eluting with an acetonitrile / 0.1 M TEAA gradient (Em. max 665 nm, H<sub>2</sub>O).


### 9. EXAMPLE: SYNTHESIS OF RHODAMINE DYE 234



Rhodamine dye **234** was synthesized as illustrated in Schemes (I) and (II) from the appropriate aminophenol starting materials.

5

#### 9.1 Synthesis of Pyrrolidinyl Phenol 214




214

10 A solution of *m*-aminophenol (12.6 g, 115 mmol) was heated in 1,4-dibromobutane (50 g, 230 mmol) at 130 °C for 18 hr. The mixtures was cooled to room temperature and the gum was triturated with ether and then ethyl acetate. The gum was dissolved in 120 ml of 1 M NaOH and extracted with ethyl acetate (100 ml). The layers were separated and the organic phase was washed with water twice and then brine. After drying over magnesium sulfate, the crude product was  
15 purified by silica gel chromatography (DCM/methanol 100:1) to give a pale yellow solid. The

solid was refluxed in toluene (500 ml) and triethylamine (16 ml, 115 mmol) for 2 hr., cooled to room temperature, and washed with water. The solvent was evaporated to leave pyrrolidinyl phenol **214** as a white solid (11 g, 60%). <sup>1</sup>H NMR (**214**, d6-DMSO) δ 8.98, 1H, s; 6.90, 1H, t; 5.95, 3H, m; 3.18, 4H, m; 1.95, 4H, m.

5

## 9.2 Synthesis of Pyrrolidinyl Ketone 216


**216**

10

A solution of aminophenol **214** (0.74 g, 4.6 mmol) and tetrafluorophthalic anhydride (1 g, 4.6 mmol) was refluxed in toluene (5 ml) for 3 hr. The solution was cooled to rt and the precipitated collected to yield pyrrolidinyl ketone **216** (1.3 g, 77%).

15

## 9.3 Synthesis of Tetrafluoro 218

**218**


5

Phosphorous oxytrichloride (1 ml, 10 mmol) was added to a solution of pyrrolidinyl ketone **216** (1.3 g, 3.5 mmol) in chloroform (10 ml). The solution was stirred for 15 min., aminophenol **214** (0.56 g, 3.5 mmol) was added and the solution was refluxed for 4 hr. The solution was cooled to rt and the reaction quenched with water (0.5 ml). Tetrafluoro **218** was purified by C18 reverse phase  
 10 chromatography (MeOH / 0.1 M TEAA, 4:1) to afford metallic green solid (0.73 g, 41%, Abs.max 576 nm, Em. max 594 nm, CH<sub>3</sub>OH).

#### 9.4 Synthesis of Rhodamine Dye 234


Rhodamine dye **234** (Abs. max 590 nm, Em. max 606 nm, CH<sub>3</sub>OH; Em. max 611  
 15 nm, H<sub>2</sub>O) was synthesized from **218** as described in Section 7.2, *supra*, and purified by C-8 reverse phase HPLC, eluting with an acetonitrile / 0.1 M TEAA gradient.

#### 10. EXAMPLE: SYNTHESIS OF RHODAMINE DYE 236



Rhodamine dye 236 was synthesized from 3-(bisbenzylamino)phenol and 5 tetrafluorophthalic anhydride 114 as illustrated in Schemes I and II.

#### 10.1 Synthesis of bis-benzyl tetrafluoro ketone 222




222

10 A solution of 3-(bisbenzylamino)phenol (14.5 g, 50 mmol) and tetrafluorophthalic anhydride (11 g, 50 mmol) were refluxed in toluene (50 ml) for 18 hr. The solvent was evaporated

and the residue purified by C18 reverse phase chromatography (MeOH/water, 7:3) to afford bis-benzyl tetrafluoro ketone **222** as a white solid (6.5 g, 25%).

## 10.2 Synthesis of Tetra-benzyl Tetrafluoro **224**




**224**

Phosphorous oxytrichloride (3.3 ml, 35 mmol) was added to a solution of bis-benzyl ketone **222** (6.0 g, 12 mmol) in chloroform (60 ml). The solution was stirred for 15 min., 3-(bisbenzylamino)phenol (3.4 g, 12 mmol) was added and the mixture was refluxed for 3 hr. The solution was cooled to room temperature and the reaction was quenched with water (1 ml). The solvent was evaporated and the residue purified by C18 reverse phase chromatography (MeOH / 0.1 M TEAA, 9:1) to afford tetra-benzyl tetrafluoro dye **224** as a dark red solid (0.62 g, 7%, Abs. max 566 nm).

## 10.3 Synthesis of Tetrafluoro **226**




**226**

A suspension of dye **224** (0.62 g, 0.8 mmol) was heated in conc. HBr (20 ml) to 110°C for 45 min. The reaction mixture was poured into ice-water and the precipitated was collected and purified by C18 reverse phase chromatography (CH<sub>3</sub>OH : 0.1M TEAA, 3:2) to afford debenzylated dye **226** as a metallic green solid (100 mg, 31%, Abs. max 521 nm).

#### 10.4 Synthesis of Rhodamine Dye 236

Rhodamine dye **236** was synthesized from **226** as described in Section 7.2, *supra*. Purification of **236** was accomplished on C18 silica gel with stepwise elution with 20-70% methanol vs. 0.1 M triethylammonium acetate. The dye **236** eluted between 30% and 50% methanol. The solvent was evaporated and the residual blue oil was precipitated with ether to provide the title dye as a blue solid (85 mg, Em. max 545 nm, H<sub>2</sub>O).

11. **EXAMPLE: SYNTHESIS OF FRET DYE 230**

230

5

11.1 **Synthesis of Succinimidyl (NHS) Ester of 196**

To a solution of dye **196** (5 mg) in DMF (100  $\mu\text{L}$ ) was added succinimidyl tetramethyluronium tetrafluoroborate (20 mg) and diisopropylamine (10  $\mu\text{L}$ ). TLC analysis on C2-reverse phase silica gel eluting with 1:1 methanol: 0.1 M triethylammonium acetate could

10 distinguish dye **196** ( $R_f = 0.2$ ) from **196** succinimidyl ester ( $R_f = 0$ ). After 1 hr, the reaction


appeared to be complete and was partitioned between 5% HCl and dichloromethane. The organic layer was dried over  $\text{Na}_2\text{SO}_4$  and the solvent evaporated to yield the **196** NHS ester as a blue solid.

### 11.2 Synthesis of FRET Dye 230

5

Energy-transfer (FRET) dye **230** is prepared by coupling **196** NHS ester and 4-aminomethylbenzoic acid and 4'-aminomethyl-6-carboxyfluorescein (Molecular Probes Inc., Eugene, OR), according to methods described in U.S. Patent No. 5,863,727. For example, 1  $\mu\text{mole}$  of **196** NHS dissolved in 250  $\mu\text{l}$  of DMSO is added to a solution of 2  $\mu\text{mole}$  4'-

10 aminomethyl-6-carboxyfluorescein in 100  $\mu\text{l}$  DMSO and 20  $\mu\text{l}$  triethylamine. After mixing the solution is let stand for about 12 hours, monitoring the progress of coupling by reverse phase HPLC.

12. **EXAMPLE: SYNTHESIS OF FRET DYE 230-ddATP**

5

FRET dye **230** is activated as the NHS ester by the method of Example *supra* 11.1 and coupled to 7-deaza-7-aminopropargyl, 2'-3' dideoxyadenosine-5'-triphosphate according to the methods in U.S. Patent No. 5,821,356 and 5,770,716 to give FRET dye **230**-ddATP.

Alternatively, 4'-trifluoroacetamidomethyl 6-carboxyfluorescein NHS ester is coupled with 7-deaza-7-aminopropargyl, 2'-3' dideoxyadenosine-5'-triphosphate to give the intermediate dye-nucleotide. The trifluoroacetyl protecting group is removed under basic conditions and the intermediate dye nucleotide is coupled with **230** NHS ester to give FRET dye **230**-ddATP by a

two-step method of conjugating the energy-transfer dye to a nucleotide.

**13. EXAMPLE: SANGER-TYPE SEQUENCING USING AN ENERGY TRANSFER DYE**

5 Following the methods described in U.S. Patents No. 5,821,356 and 5,366,860, Sanger-type terminator sequencing was performed on pGEM (PE Biosystems, Foster City, CA) using Taq FS polymerase (PE Biosystems, Foster City, CA), a mixture of four dNTPs, an unlabeled sequencing primer and a single labeled terminator (6-FAM-230-ddATP). A plot of the resultant sequencing data, obtained on an ABI PRISM Model 310 instrument (PE Biosystems, Foster City,  
10 CA) is provided in FIG. 1.

**14. EXAMPLE: FOUR-COLOR SANGER-TYPE SEQUENCING**

Following the methods described in U.S. Patents No. 5,821,356 and 5,366,860, four-color Sanger-type terminator sequencing was performed on pGEM1 (PE Biosystems, Foster City, CA)  
15 using Taq FS polymerase (PE Biosystems, Foster City, CA), a mixture of four dNTPs, an unlabeled sequencing primer and a mixture of four, 3'-fluoro, spectrally-resolvable labeled terminators (6-FAM-230-ddATP; 5-FAM-236-7-deaza-ddGTP; 6-FAM-JON-ddTTP; 6-FAM-ROX-ddCTP). A plot of the resultant sequencing data, obtained on an ABI PRISM Model 310 instrument (PE Biosystems, Foster City, CA) is provided in FIG. 2.

20 As can be seen in FIG. 2, all of the dye-labeled polynucleotides exhibit significant fluorescence intensity. Moreover, the different dye-labeled polynucleotide exhibit sufficiently similar mobilities, resulting in good resolution.

**15. EXAMPLE: ANTI-HUMAN ANTIBODY-DYE CONJUGATE DETECTION**

**25 Anti-human IL-8 antibody-dye - 196 conjugate**

Polyclonal anti-human IL-8 antibody (R&D Systems, Minneapolis, MN; 0.5 mg in 0.5 ml PBS) was incubated with 50  $\mu$ l of 1 M  $\text{Na}_2\text{CO}_3$  and 105  $\mu$ g of dye 196 for 1 hour at room temperature, in the dark. The antibody-dye - 196 conjugate (Abs. max 666 nm,  $\text{H}_2\text{O}$ ) was separated from the free dye using a sephadex G-50 (fine) size exclusion column. The conjugate  
30 was passed through a 0.2  $\mu$  filter and stored at 4° C in PBS. The concentration of the antibody was determined according to the following equation:  $[A_{280} - (0.82 \times A_{640})] / 170,000$ , where 0.82 =  $A_{280}/A_{640}$  of the free dye and 170,000 is the extinction coefficient for the antibody. The

concentration of the dye was determined according to the following equation:  $A_{640}/120,000$ , where 120,000 is the extinction coefficient for dye 196. The dye 196-labeled polyclonal anti-human IL-8 antibody concentration was 0.18 mg/ml with an F/P ratio of 2.3/1.

5 Preparation of monoclonal anti-human IL-8 antibody coated beads

Goat anti-mouse IgG (Fc) polystyrene beads (200 ml of 0.5% w/v; 6  $\mu$ m bead diameter; Spherotech) were first washed 3 times by centrifugation and resuspension of the bead pellet with 1 ml PBS. 4  $\mu$ g of monoclonal anti-human IL-8 antibodies (R&D Systems) was added to the bead suspension for a final volume of 1 ml. After incubation for 16 hrs at room temperature with gentle  
10 mixing, the monoclonal antibody-coated beads were washed 2 times as described above and resuspended in 1 ml PBS. The final bead concentration was 0.1% w/v or  $8 \times 10^6$  beads/ml. The monoclonal antibody-coated beads were stable for approximately 1 month when stored at 4°C.

Fluorescent-linked Immunosorbent Assay (FLISA)

15 To generate a standard curve, a two-fold serial dilution of human IL-8 peptide (R&D Systems) in 50  $\mu$ l FLISA buffer (PBS containing 1 mg/ml BSA, 0.35 M NaCl, 0.1% Tween 20, and 0.01%  $\text{NaN}_3$ ) were aliquoted into a 96-well plate (Corning Costar No. 3904). Subsequently, 50  $\mu$ l of a mixture containing monoclonal anti-human IL-8 antibody-coated beads and dye 196-labeled polyclonal anti-human IL-8 antibodies in FLISA buffer was then added to each well. The final  
20 concentrations in a total volume of 100  $\mu$ l per well were 64,000 antibody-coated beads/ml and 0.11  $\mu$ g/ml of dye 196-labeled polyclonal anti-human IL-8 antibodies. The serial dilution for IL-8 peptide ranged from 2,000-2.0 pg/ml. The assay contained eight replicates, and control wells containing no peptide were included. After incubating overnight at room temperature in the dark, the 96-well plate was scanned using the FMAT 8100 HTS System (PE Biosystems, Foster City,  
25 CA), a macroconfocal imaging system equipped with a 633-nm HeNe laser and 2-channel fluorescence detection (FL1: 650-685 nm, FL2: 685-720 nm). A 1 mm<sup>2</sup> area of each well is raster scanned at a depth of focus of 100  $\mu$ m and at a rate of 1 sec/well, with 256 scan lines generated. The fluorescence intensity associated with each bead is then obtained from the digitized images and the average fluorescence intensity per bead is calculated per well. A log vs log graph of the  
30 average fluorescence intensity in FL1 vs pg/ml of the IL-8 peptide is shown in Figure 4. The linear dynamic range of the assay is 250-3.9 pg/ml IL-8.

While the present invention has been described by reference to the preferred embodiments and examples detailed above, it is to be understood that these examples are intended in an illustrative, rather than limiting, sense. It is contemplated that modifications that do not depart from the spirit of the invention will readily occur to those of skill in the art. Any such  
5 modifications are intended to fall within the scope of the appended claims.

All of the references, publications and patents cited in the specification are incorporated herein by reference to the same extent as if each reference were individually incorporated by reference.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207